

Partial Applicative Theories and Explicit Substitutions

Thomas Strahm*

Institut für Informatik und angewandte Mathematik, Universität Bern
Länggassstrasse 51, CH-3012 Bern, Switzerland
(strahm@iam.unibe.ch)

May 15, 1996

Abstract

Systems based on theories with partial self-application are relevant to the formalization of constructive mathematics and as a logical basis for functional programming languages. In the literature they are either presented in the form of partial combinatory logic or the partial λ calculus, and sometimes these two approaches are erroneously considered to be equivalent.

In this paper we address some defects of the partial λ calculus as a constructive framework for partial functions. In particular, the partial λ calculus is not embeddable into partial combinatory logic and it lacks the standard recursion-theoretic model. The main reason is a concept of substitution, which is not consistent with a strongly intensional point of view. We design a weakening of the partial λ calculus, which can be embedded into partial combinatory logic. As a consequence, the natural numbers with partial recursive function application *are* a model of our system. The novel point will be the use of *explicit substitutions*, which have previously been studied in the literature in connection with the implementation of functional programming languages.

Keywords: Explicit mathematics, logic of partial terms, partial λ calculus, partial combinatory logic, explicit substitutions

1 Introduction

Partial applicative theories form the basis of various formal systems for constructive mathematics and functional programming. Feferman introduced in [6] and [7] partial applicative theories of operations and classes in order to give a logical account to

*Research supported by the Swiss National Science Foundation.

Bishop’s style of constructive mathematics (BCM). More recently, Feferman’s systems of explicit mathematics were used to develop a unitary axiomatic framework for representing programs, stating properties of programs, and proving properties of programs. The programs considered are taken from functional programming languages, which are mainly based on the untyped λ calculus. Important references for the use of systems of explicit mathematics in the context of functional programming are Feferman [8], [9], [10], Jäger [16], [17] and Marzetta [22], [23].

As far as the explicit representation of a theory with partial self-application in the previous literature is concerned, people either took partial combinatory logic or the partial λ calculus (without the rule ξ) as the applicative basis. The former possibility was chosen in [2], [6], [7], [11], [16], [17], [23], [29], the latter in [8], [9], [10] and [22]. At first sight, these two approaches seem to be completely equivalent, and sometimes they are treated as such in the literature. But they are only equivalent in the presence of a *total* logic, since then λ calculus (without ξ) can be embedded into combinatory logic and vice versa. For a detailed discussion of the total case also in the context of reductions, the reader is referred to Hindley [14], and Hindley and Longo [15].

The situation changes drastically if one considers a *partial* application operation. Then the partial λ calculus (without ξ) is no longer embeddable into partial combinatory logic. This is due to the fact that the coding of λ abstraction in the context of partial combinatory logic is more complicated than usual. The modified definition of λ does not permit to push a substitution θ inside an abstraction $(\lambda x.t)$, a principle, which is valid for the partial λ calculus for obvious reasons. For example, the terms $(\lambda x.y)[zz/y]$ and $(\lambda x.zz)$ are not equal in partial combinatory logic. Hence, the stronger concept of substitution in the partial λ calculus makes its embedding into partial combinatory logic fail. For the same reason, the standard recursion-theoretic model of partial combinatory logic is no longer a model of the partial λ calculus. Recently, Pezzoli [26] even proved that there is no (reasonable) recursion-theoretic interpretation of the partial λ calculus at all.

As a consequence, it is not possible to determine proof-theoretical upper bounds of applicative theories based on the partial λ calculus by means of the recursion-theoretic model, as it can be done for the corresponding systems based on partial combinatory logic. Although the upper bounds of those systems can be determined by formalizing a *total* term model (cf. Jäger and Strahm [18]), the adequacy of the partial λ calculus as a constructive framework for partial functions is seriously put into question. The system simply does not seem to have any reasonable models with a perspicuous constructive meaning that are truly partial. Not only the recursion-theoretic model but also other partial models of partial combinatory logic do not have their counterparts as models of the partial λ calculus.

In the following we propose a modification of the partial λ calculus, which can be embedded into partial combinatory logic via a natural embedding. As a consequence, this weakened form of the partial λ calculus has all the partial models which we have for partial combinatory logic. In particular, it is possible to determine upper bounds of systems of explicit mathematics based on our modified version of the partial λ calculus using the recursion-theoretic model.

The novel point of our system will be the use of *explicit substitutions*. According to this approach substitution is no longer a notion of the metalanguage, but an operation axiomatized in the theory under consideration. If t, s_1, \dots, s_n are terms and θ is the substitution $\{s_1/x_1, \dots, s_n/x_n\}$ then $t\theta$ is no longer an abbreviation in the metalanguage for the term t with the variables x_i simultaneously replaced by the terms s_i , but a purely syntactical object. The evaluation of θ has to be described by appropriate axioms or rules. So it is possible to provide a very controlled process of substitution. In particular, substitution can be axiomatized in a way that is consistent with the recursion-theoretic model and partial combinatory logic, respectively. Hence, a substitution θ can no longer be pushed inside an abstraction $(\lambda x.t)$.

The theory of explicit substitutions has been treated in the literature before, but from a different point of view. The main work has been done in the context of implementation of functional programming languages, and application in those systems is always total. The very concern of the present work, however, is to study a *partial* application operation. A key reference for the previous work on explicit substitution is the paper by Abadi, Cardelli, Curien and Lévy [1]. Further investigations are presented in Curien [4], Curien, Hardin and Lévy [5], Hardin and Lévy [13] as well as in Lescanne and Rouyer [20].

Recently, Martin-Löf [21] introduced a calculus of explicit substitutions in connection with his intuitionistic theory of types, which is worked out in Tasistro [28].

Let us briefly sketch the procedure of these investigations. In Section 2 we first introduce partial combinatory logic \mathbf{CL}_p and the (usual) partial λ calculus λ_p (without ξ). In particular, we recapitulate Beeson's logic of partial terms. After having sketched some interesting partial models of \mathbf{CL}_p , we discuss the substitution problems, which prevent the embedding of λ_p into \mathbf{CL}_p , and we give Pezzoli's result mentioned above. In Section 3 we give a detailed formulation of the system $\lambda_p\sigma$, which is a modification of λ_p by explicit substitutions. The system incorporates an adaptation of Beeson's logic of partial terms to the framework of explicit substitutions, and rules to evaluate substitutions, of course. We further show that $\lambda_p\sigma$ is embeddable into \mathbf{CL}_p via a natural embedding and that \mathbf{CL}_p is also contained in $\lambda_p\sigma$ via the standard embedding. In Section 4, finally, we study the reduction relation on $\lambda_p\sigma$ terms corresponding to the system $\lambda_p\sigma$. We give a long and tedious proof

for the Church Rosser property of this relation.

We finish this introduction by mentioning that recently Stärk [27] has established a natural relationship between the $\lambda_p\sigma$ calculus and the programming language SCHEME.

2 The systems \mathbf{CL}_p and λ_p

Let us first define partial combinatory logic \mathbf{CL}_p and the usual partial λ calculus λ_p , without the rule ξ . The language of \mathbf{CL}_p includes an infinite list of object variables (in the metalanguage: $x, y, z, f, g, h, u, v, w, \dots$), constants \mathbf{k} and \mathbf{s} (partial combinatory algebra), the binary function symbol \circ (application), the equality symbol $=$, the symbol \downarrow (defined) and the usual propositional connectives and first order quantifiers. The language of λ_p contains the same symbols except that \mathbf{k} and \mathbf{s} are replaced by the abstractor λ .

The terms of \mathbf{CL}_p and λ_p (in the metalanguage: r, s, t, \dots) are given by the following definitions.

Definition 1 (\mathbf{CL}_p terms)

1. Every variable is a \mathbf{CL}_p term.
2. \mathbf{k} and \mathbf{s} are \mathbf{CL}_p terms.
3. If s and t are \mathbf{CL}_p terms, then $(s \circ t)$ is a \mathbf{CL}_p term.

Definition 2 (λ_p terms)

1. Every variable is a λ_p term.
2. If t is a λ_p term, then $(\lambda x.t)$ is a λ_p term.
3. If s and t are λ_p terms, then $(s \circ t)$ is a λ_p term.

In the following we write (st) for $(s \circ t)$. Additionally, we adopt the convention of association to the left, i.e. $t_1 t_2 t_3 \dots t_n$ stands for $(\dots((t_1 t_2) t_3) \dots t_n)$. Finally, we often write $(\lambda x_1 \dots x_n.t)$ instead of $\lambda x_1.(\lambda x_2.(\dots(\lambda x_n.t)\dots))$.

The formulas of \mathbf{CL}_p (in the metalanguage: A, B, C, \dots) are defined in the obvious way.

Definition 3 (\mathbf{CL}_p formulas)

1. If s and t are \mathbf{CL}_p terms, then $(s = t)$ is a \mathbf{CL}_p formula.

2. If t is a \mathbf{CL}_p term, then $t \downarrow$ is a \mathbf{CL}_p formula.
3. If A and B are \mathbf{CL}_p formulas, then $\neg A$, $(A \vee B)$, $(A \wedge B)$ and $(A \rightarrow B)$ are \mathbf{CL}_p formulas.
4. If A is a \mathbf{CL}_p formula, then $\exists xA$ and $\forall xA$ are \mathbf{CL}_p formulas.

The formulas of λ_p are defined in exactly the same way as the \mathbf{CL}_p formulas.

If t is a term of \mathbf{CL}_p or λ_p , then $fvar(t)$ denotes the set of free variables of t . As usual, $t[s_1/x_1, \dots, s_n/x_n]$ is the term t in which the free occurrences of x_1, \dots, x_n are simultaneously substituted by s_1, \dots, s_n . If t is a λ_p term, then in the case of variable collisions bound variables have to be renamed. Analogously, $fvar(A)$ and $A[s_1/x_1, \dots, s_n/x_n]$ are defined.

Definition 4

1. $(A \leftrightarrow B) := ((A \rightarrow B) \wedge (B \rightarrow A))$.
2. $(t \simeq s) := ((t \downarrow \vee s \downarrow) \rightarrow t = s)$.

Together with the axioms stated below it will become clear that $(t \simeq s)$ is equivalent to

$$(t \downarrow \wedge s \downarrow \wedge t = s) \vee (\neg t \downarrow \wedge \neg s \downarrow),$$

i.e. \simeq is a partial equality relation as it is used e.g. in a recursion-theoretic framework.

In the following we give the axioms and rules of inference of the systems \mathbf{CL}_p and λ_p , respectively. The logic of both \mathbf{CL}_p and λ_p is the classical logic of partial terms due to Beeson [3] and [2]. It corresponds to \mathbf{E}^+ logic as it is discussed in Troelstra and van Dalen [30]. All our results also hold if *intuitionistic* logic is chosen as a basis of \mathbf{CL}_p and λ_p , respectively.

Definition 5 The system \mathbf{CL}_p is formulated in the language of \mathbf{CL}_p , and contains the following list of axioms and rules of inference.

A. PROPOSITIONAL LOGIC

- (1) Some complete axiom schemes of classical propositional logic

B. QUANTIFIER AXIOMS

- (2) $\forall xA \wedge t \downarrow \rightarrow A[t/x]$
- (3) $A[t/x] \wedge t \downarrow \rightarrow \exists xA$

C. EQUALITY AXIOMS

- (4) $x = x \quad t = s \rightarrow s = t \quad t = s \wedge s = r \rightarrow t = r$
- (5) $t_1 = s_1 \wedge t_2 = s_2 \rightarrow t_1 t_2 \simeq s_1 s_2$

D. STRICTNESS AXIOMS

$$(6) \ x \downarrow$$

$$(7) \ t_1 = t_2 \rightarrow t_1 \downarrow \wedge t_2 \downarrow \quad t_1 t_2 \downarrow \rightarrow t_1 \downarrow \wedge t_2 \downarrow$$

$$(8) \ \mathbf{k} \downarrow \quad \mathbf{s} \downarrow \quad \mathbf{s}xy \downarrow$$

E. PARTIAL COMBINATORY ALGEBRA

$$(9) \ \mathbf{k}xy \simeq x$$

$$(10) \ \mathbf{s}xyz \simeq xz(yz)$$

F. RULES OF INFERENCE

$$(11) \ \frac{A \quad A \rightarrow B}{B}$$

$$(12) \ \frac{A \rightarrow B}{\exists x A \rightarrow B} \quad \frac{A \rightarrow B}{A \rightarrow \forall x B}$$

In the inference rules (12) x does not appear free in the conclusion.

Definition 6 The system λ_p is formulated in the language of λ_p and contains the same axioms and rules of inference as \mathbf{CL}_p except that the strictness axioms (8) are replaced by

$$(\lambda x.t) \downarrow$$

and the axioms of a partial combinatory algebra are replaced by the β axiom

$$(\beta) \ (\lambda x.t)y \simeq t[y/x].$$

It is important to notice that in the system λ_p we do not have the partial analogue of the rule (ξ) ,

$$(\xi) \ \frac{t \simeq s}{\lambda x.t = \lambda x.s}.$$

The principle (ξ) induces a weak form of extensionality, which is not consistent with the strongly intensional character of (indices of) partial functions as met e.g. in a recursion-theoretic framework. Formally, this means that it is a priori hopeless to embed λ calculus into combinatory logic in the presence of (ξ) . Moreover, the absence of (ξ) is in accord with most implementations of λ calculus based languages: Functions are considered as values, and are only evaluated when arguments are fed in. As already mentioned in the introduction, λ calculus without (ξ) in the context of a *total* application operation is discussed in [14] and [15].

Nevertheless, we will shortly address fully extensional versions of \mathbf{CL}_p and λ_p , respectively, i.e. we will consider the strong extensionality axiom (Ext),

$$\text{(Ext)} \quad \forall x (fx \simeq gx) \rightarrow (f = g).$$

Let us briefly sketch some models of \mathbf{CL}_p . As we are mainly interested in partiality, we will only discuss truly partial models. Of course, there are many models of \mathbf{CL}_p where application is a total operation, e.g. each model of the λ calculus is a model of \mathbf{CL}_p . In the following we give two partial models of \mathbf{CL}_p .

The recursion-theoretic model PRO. The universe of the model *PRO* of partial recursive operations consists of the set of natural numbers ω . Application \circ is interpreted as partial recursive function application, i.e. $x \circ y$ means $\{x\}(y)$ in *PRO*, where $\{x\}$ is a standard enumeration of the partial recursive functions. It is easy to find appropriate interpretations of \mathbf{k} and \mathbf{s} such that the axioms of a partial combinatory algebra are satisfied. *PRO* provides a natural example of a domain where objects may be programs as well as inputs to programs. The model underlines the constructive and operational character of applicative theories.

The normal term model CNT. This model is based on standard notions of term reduction for combinatory logic, i.e. $\mathbf{k}t_1t_2$ reduces to t_1 and $\mathbf{s}t_1t_2t_3$ reduces to $t_1t_3(t_2t_3)$. The universe of *CNT* consists of all closed \mathbf{CL}_p terms in normal form, \mathbf{k} and \mathbf{s} are interpreted by themselves, and $t_1 \circ t_2$ means $\text{InFirst}(t_1t_2)$. $\text{InFirst}(t_1t_2)$ denotes the uniquely determined normal term s provided that t_1t_2 can be reduced to s according to the *leftmost minimal strategy*, $\text{InFirst}(t_1t_2)$ is undefined otherwise. Using the leftmost minimal strategy, at each stage of a reduction sequence the leftmost minimal redex is contracted, where a redex is called *minimal*, if it does not contain any other redexes. It is necessary to use the leftmost minimal strategy in order to be consistent with the strictness axioms of \mathbf{CL}_p . The model *CNT* provides us with another interesting operational semantics of \mathbf{CL}_p . For a detailed description of *CNT* the reader is referred to Beeson [2], p. 119 ff.

As we will see below, the models just described cannot be made into models of the system λ_p in a reasonable way. This is due to a stronger concept of substitution, which is inherent in the partial λ calculus λ_p .

Our next aim is to code λ abstraction in \mathbf{CL}_p . We have to be careful in defining it in the context of the logic of partial terms, because we want $\lambda x.t$ to be defined for each term t . As we will see below, this modified λ abstraction will have very unpleasant properties as far as substitution is concerned.

Definition 7 ($\lambda^*x.t$)

For each term t of \mathbf{CL}_p a term $\lambda^*x.t$ is defined by induction on the complexity of t .

1. If t is the variable x , then $\lambda^*x.t := \mathbf{s}kk$.
2. If t is a variable different from x or a constant, then $\lambda^*x.t := \mathbf{k}t$.
3. If t is the term (t_1t_2) , then $\lambda^*x.t := \mathbf{s}(\lambda^*x.t_1)(\lambda^*x.t_2)$.

Lemma 8 *We have for all \mathbf{CL}_p terms t and s :*

1. $fvar(\lambda^*x.t) = fvar(t) \setminus \{x\}$.
2. $\mathbf{CL}_p \vdash \lambda^*x.t \downarrow$.
3. $\mathbf{CL}_p \vdash (\lambda^*x.t)x \simeq t$.
4. $\mathbf{CL}_p \vdash s \downarrow \rightarrow (\lambda^*x.t)s \simeq t[s/x]$.

PROOF (1)–(3) are proved by induction on the complexity of t . (4) is a direct consequence of (3). \square

In the context of a total logic, one normally defines $\lambda^*x.t := \mathbf{k}t$, if $x \notin fvar(t)$. So we have e.g. $\lambda^*x.(yz) = \mathbf{k}(yz)$. The example shows that $\lambda^*x.t \downarrow$ does not hold for the usual definition of λ abstraction.

As already mentioned, the λ abstraction of Definition 7 behaves very badly as far as substitution in λ expressions is concerned. For a usual λ abstraction we have

$$(\lambda x.t)[s/y] = \lambda x.t[s/y], \quad (*)$$

provided that $x \neq y$ and $x \notin fvar(s)$. This property (which we only need for $s \downarrow$ in a partial setting) fails for the λ^* defined above: We have e.g. $(\lambda^*x.y)[zz/y] = \mathbf{k}(zz)$, but $\lambda^*x.zz = \mathbf{s}(\mathbf{k}z)(\mathbf{k}z)$.

The fact that the substitution property (*) does not hold for the λ abstraction of Definition 7 is not just a technical inconvenience, but has rather strong consequences for the system λ_p as a constructive framework for partial functions. Since (*) trivially holds in λ_p , the standard embedding of λ_p into \mathbf{CL}_p fails. As a consequence, the \mathbf{CL}_p models *PRO* and *CNT* described above do not translate into models of λ_p .

An illustrative consequence of the stronger substitution concept of λ_p is a very weak form of (ξ), which is derivable in λ_p . Let s, t be λ_p terms, and x be a variable with $x \notin fvar(s) \cup fvar(t)$. Then it is easy to see that the principle

$$s = t \rightarrow \lambda x.s = \lambda x.t \quad (**)$$

is derivable in λ_p *only* from (β) and the fact that equality respects application. For $(s = t)$ implies $(\lambda y.\lambda x.y)s \simeq (\lambda y.\lambda x.y)t$, and hence we can conclude by the β axiom

$$\lambda x.s = (\lambda y.\lambda x.y)s = (\lambda y.\lambda x.y)t = \lambda x.t.$$

Note that in the argument above, we pushed the substitutions $[s/y]$ and $[t/y]$ inside the abstraction $(\lambda x.y)$. The principle $(**)$ can be considered as an extremely weak form of extensionality, and therefore, it has to be rejected from a strongly intensional point of view.

We have seen above that the λ abstraction of Definition 7 does not yield a recursion-theoretic interpretation of λ_p . The question arises whether it is possible to find another encoding of λ in the model PRO satisfying the substitution property $(*)$. It is easily checked that all λ encodings known from the literature (cf. e.g. Kleene [19], p. 344 ff.) do not validate $(*)$, and all attempts to define another such λ failed. Very recently, Pezzoli [26] found an elegant formal argument showing that the existence of a recursion-theoretic interpretation of λ_p which has a partial recursive term evaluation function contradicts the undecidability of the halting problem.

Theorem 9 (Pezzoli) *It is not possible to make the partial recursive functions model PRO of \mathbf{CL}_p into a model of the partial λ calculus λ_p in such a way that the term evaluation function $f(t, \rho) \simeq \|t\|_\rho$, for t a λ_p term and ρ an assignment of variables, is partial recursive.*

PROOF Let us assume that we can make the partial recursive functions into a model of λ_p such that $\|t\|_\rho$ is partial recursive. By (4) of Lemma 8 and the substitution property $(*)$ we know that λ_p proves

$$(zz) \downarrow \rightarrow (\lambda xy.x)(zz) = \lambda y.zz,$$

which immediately yields

$$(zz) \downarrow \rightarrow \lambda w.((\lambda xy.x)(zz)) = \lambda wy.zz \quad (***)$$

by the principle $(**)$ mentioned above. Now consider $\|\lambda w.((\lambda xy.x)(zz))\|_\rho = a_\rho$ and $\|\lambda wy.zz\|_\rho = b_\rho$, which are defined for every ρ since λ abstraction is always defined. By $(***)$ we have $a_\rho = b_\rho$ whenever $\{\rho(z)\}(\rho(z)) \downarrow$; if $\{\rho(z)\}(\rho(z)) \uparrow$, then a_ρ must be an index of the always undefined function and b_ρ an index for the constant function $n \mapsto \|\lambda y.zz\|_\rho$, so in this case $a_\rho \neq b_\rho$. However, by hypothesis, we can compute a_ρ and b_ρ , and therefore, we can decide whether $\{n\}(n) \downarrow$. This is not possible. \square

The following corollary is immediate from the fact that PRO is a model of \mathbf{CL}_p .

Corollary 10 *There is no recursive encoding of λ in \mathbf{CL}_p validating $(*)$.*

It should be stressed that the problems described above completely disappear in the presence of the extensionality axiom (Ext). In particular, $(*)$ holds in $\mathbf{CL}_p + (\text{Ext})$.

Lemma 11 *We have for all \mathbf{CL}_p terms t, s and all variables x, y such that $x \neq y$ and $x \notin \text{fvar}(s)$:*

$$\mathbf{CL}_p + (\text{Ext}) \vdash s \downarrow \rightarrow (\lambda^*x.t)[s/y] = \lambda^*x.t[s/y].$$

PROOF Assume $s \downarrow$. By Lemma 8 we have $(\lambda^*x.t)x \simeq t$ and $(\lambda^*x.t[s/y])x \simeq t[s/y]$. Since $s \downarrow$ and $x \neq y$ we can conclude that

$$(\lambda^*x.t)[s/y]x \simeq t[s/y] \simeq (\lambda^*x.t[s/y])x,$$

which by (Ext) immediately implies the claim of the lemma. \square

As an immediate consequence of this lemma we get the following theorem, which is also stated in Moggi [24].

Theorem 12 *The systems $\mathbf{CL}_p + (\text{Ext})$ and $\lambda_p + (\text{Ext})$ are equivalent with respect to the standard embeddings.*

3 The system $\lambda_p\sigma$

In the following, we introduce the system $\lambda_p\sigma$, which is a modification of the system λ_p by explicit substitutions. The language of $\lambda_p\sigma$ is an extension of the language of λ_p by the set brackets $\{, \}$, the slash $/$ and commas. The new symbols will be used in order to form finite sets of variable bindings, i.e. *substitutions*.

The terms (in the metalanguage: r, s, t, \dots) and substitutions (in the metalanguage: $\theta, \sigma, \tau, \dots$) of $\lambda_p\sigma$ are given by a simultaneous inductive definition.

Definition 13 (*$\lambda_p\sigma$ terms and $\lambda_p\sigma$ substitutions*)

1. Every variable is a $\lambda_p\sigma$ term.
2. If t is a $\lambda_p\sigma$ term, then $(\lambda x.t)$ is a $\lambda_p\sigma$ term.
3. If s and t are $\lambda_p\sigma$ terms, then $(s \circ t)$ is a $\lambda_p\sigma$ term.
4. If t is a $\lambda_p\sigma$ term and if θ is a $\lambda_p\sigma$ substitution, then $(t\theta)$ is a $\lambda_p\sigma$ term.
5. If t_1, \dots, t_n are $\lambda_p\sigma$ terms and if x_1, \dots, x_n are variables with $x_i \neq x_j$ for $1 \leq i < j \leq n$, then $\{t_1/x_1, \dots, t_n/x_n\}$ is a $\lambda_p\sigma$ substitution.

In the following we will often write $t\theta\sigma$ instead of $(t\theta)\sigma$.

The formulas of $\lambda_p\sigma$ are defined in exactly the same way as the formulas of λ_p , e.g. we have an atomic formula $t \downarrow$ for each $\lambda_p\sigma$ term t . In the following we often

speak of terms, substitutions, and formulas instead of $\lambda_p\sigma$ terms, $\lambda_p\sigma$ substitutions, and $\lambda_p\sigma$ formulas.

We again want to stress the difference between $t[s/x]$ and $t\{s/x\}$. In the first expression substitution is an abbreviation in the metalanguage for the term t with all free occurrences of x replaced by s . $t\{s/x\}$, however, is a purely syntactical object where the substitution $\{s/x\}$ can only be evaluated by means of appropriate axioms to be described below.

We will often use the following abbreviations.

Definition 14 Let $\theta = \{t_1/x_1, \dots, t_n/x_n\}$ and $\theta' = \{s_1/x_1, \dots, s_n/x_n\}$ be substitutions and let r be an arbitrary term. Then we define:

1. $dom\theta := \{x_1, \dots, x_n\}$.
2. $\theta \downarrow := t_1 \downarrow \wedge \dots \wedge t_n \downarrow$.
3. $\theta \simeq \theta' := t_1 \simeq s_1 \wedge \dots \wedge t_n \simeq s_n$.
4. $t/x \cdot \theta := \{t/x\} \cup \theta^{-x}$, where θ^{-x} is θ with a possible binding for x deleted, i.e. $t/x \cdot \theta$ denotes the update of θ by $\{t/x\}$.
5. $\varepsilon := \{\}$ (the empty substitution).

The set of free variables $fvar(t)$ of a $\lambda_p\sigma$ term t is computed in the obvious way. For reasons of completeness we give the exact inductive definition below.

Definition 15 ($fvar(t)$)

1. If t is the variable x , then $fvar(t) := \{x\}$.
2. If t is the term $(\lambda x.s)$, then $fvar(t) := fvar(s) \setminus \{x\}$.
3. If t is the term $(t_1 t_2)$, then $fvar(t) := fvar(t_1) \cup fvar(t_2)$.
4. If t is the term $(s\theta)$ and $\theta = \{s_1/x_1, \dots, s_n/x_n\}$, then

$$fvar(t) := (fvar(s) \setminus dom\theta) \cup \bigcup_{i \in I} fvar(s_i),$$

where $I = \{i : 1 \leq i \leq n \text{ and } x_i \in fvar(s)\}$.

Once $fvar(t)$ is defined, we get $fvar(A)$ as the set of free variables of a $\lambda_p\sigma$ formula A in the usual way.

We will sometimes use the notation $\theta \upharpoonright t$ for the substitution θ with all variable bindings s/y deleted for $y \notin fvar(t)$.

In the sequel we define a $\lambda_p\sigma$ formula $A\theta$ for each $\lambda_p\sigma$ formula A and each $\lambda_p\sigma$ substitution θ . The definition is by induction on the complexity of A .

Definition 16 ($A\theta$)

1. If A is the formula $(s = t)$, then $A\theta$ is the formula $(s\theta = t\theta)$.
2. If A is the formula $t \downarrow$, then $A\theta$ is the formula $t\theta \downarrow$.
3. If A is the formula $\neg B$, $(B \vee C)$, $(B \wedge C)$ or $(B \rightarrow C)$, then $A\theta$ is the formula $\neg(B\theta)$, $(B\theta \vee C\theta)$, $(B\theta \wedge C\theta)$ or $(B\theta \rightarrow C\theta)$, respectively.
4. If A is the formula $\exists xB$ or $\forall xB$, then $A\theta$ is the formula $\exists y(B[y/x]\theta)$ or $\forall y(B[y/x]\theta)$ respectively, where y is a “fresh” variable.¹

The *composition* of two substitutions θ and σ is defined in the usual way.

Definition 17 Let $\theta = \{t_1/x_1, \dots, t_n/x_n\}$ and $\sigma = \{s_1/y_1, \dots, s_m/y_m\}$ be substitutions. The substitution $\theta\sigma$ is obtained by deleting all variable bindings of the form s_i/y_i in the set

$$\{t_1\sigma/x_1, \dots, t_n\sigma/x_n, s_1/y_1, \dots, s_m/y_m\},$$

such that $y_i = x_j$ for a $1 \leq j \leq n$.

Now we are ready to give the exact formulation of the system $\lambda_p\sigma$. The logic of $\lambda_p\sigma$ is an adaptation of Beeson’s logic of partial terms to the framework of explicit substitutions. The novel point of this axiom system compared to the system λ_p are the substitution axioms (E), which incorporate rules to evaluate substitutions step by step. Furthermore, an extended form of the β axiom in the context of explicit substitutions is given.

Definition 18 The system $\lambda_p\sigma$ is formulated in the language of $\lambda_p\sigma$ and contains the following list of axioms and rules of inference.

A. PROPOSITIONAL LOGIC

- (1) Some complete axiom schemes of classical propositional logic

B. QUANTIFIER AXIOMS

- (2) $\forall \vec{x} A \wedge \theta \downarrow \rightarrow A\theta$
- (3) $A\theta \wedge \theta \downarrow \rightarrow \exists \vec{x} A$ ($\vec{x} = x_1, \dots, x_n$; $dom\theta = \{\vec{x}\}$)

C. EQUALITY AXIOMS

- (4) $x = x$ $t = s \rightarrow s = t$ $t = s \wedge s = r \rightarrow t = r$

¹ $B[y/x]$ is the formula B , where each free occurrence of x is replaced by y in the *usual* sense. The exact definition of $B[y/x]$ is straightforward, but tedious.

$$(5) \quad t_1 = s_1 \wedge t_2 = s_2 \rightarrow t_1 t_2 \simeq s_1 s_2$$

$$(6) \quad \theta \simeq \theta' \rightarrow t\theta \simeq t\theta'$$

D. STRICTNESS AXIOMS

$$(7) \quad x \downarrow$$

$$(8) \quad t_1 = t_2 \rightarrow t_1 \downarrow \wedge t_2 \downarrow \quad t_1 t_2 \downarrow \rightarrow t_1 \downarrow \wedge t_2 \downarrow$$

$$(9) \quad (\lambda x.t) \downarrow$$

E. SUBSTITUTION AXIOMS

$$(10) \quad x\theta \simeq t \quad (t/x \in \theta)$$

$$(11) \quad (ts)\theta \simeq (t\theta)(s\theta)$$

$$(12) \quad (t\theta)\sigma \simeq t(\theta\sigma)$$

$$(13) \quad t(s/x \cdot \theta) \simeq t\theta \quad (x \notin fvar(t) \cup dom\theta)$$

$$(14) \quad t\varepsilon \simeq t$$

F. β AXIOM

$$(15) \quad (\lambda x.t)\theta y \simeq t(y/x \cdot \theta)$$

G. RULES OF INFERENCE

$$(16) \quad \frac{A \quad A \rightarrow B}{B}$$

$$(17) \quad \frac{A \rightarrow B}{\exists x A \rightarrow B} \quad \frac{A \rightarrow B}{A \rightarrow \forall x B}$$

In the inference rules (17) x does not appear free in the conclusion.

It should be observed that among the substitution axioms (E) we do *not* have an axiom, which allows us to push a substitution θ inside an abstraction $(\lambda x.t)$. This is exactly what we want to prevent. Terms of the form $(\lambda x.t)\theta$ can only be resolved if applied to another object, say y . This is reflected in the extended β axiom (15), where an interleaving substitution θ is allowed. If θ is the empty substitution ε then we have the usual β axiom.

Weak λ calculi (i.e. λ calculi without (ξ) or substitution under λ) with explicit substitutions have been considered in the literature before. These include Curien's $\lambda\rho$ calculus [4], the conditional weak theory $\lambda\sigma_{cw}$ in Curien, Hardin and Lévy [5] as well as the weak theory $\lambda\sigma_w$ of [5]. The main difference (among other minor

differences) between $\lambda\rho$, $\lambda\sigma_{cw}$, $\lambda\sigma_w$ and $\lambda_p\sigma$ lies in the fact that application in the former three calculi is always *total*, whereas the main concern of the $\lambda_p\sigma$ calculus is to model a *partial* application operation. As we argued in the previous section, such a partial λ calculus must not allow (ξ) and substitution under λ in order to be consistent with the intended recursion-theoretic interpretation.

Lemma 19 *We have for all terms t, s and all substitutions θ :*

1. $\lambda_p\sigma \vdash s \downarrow \rightarrow (\lambda x.t)s \simeq t\{s/x\}$.
2. $\lambda_p\sigma \vdash s \downarrow \rightarrow (\lambda x.t)\theta s \simeq t(s/x \cdot \theta)$.
3. $\lambda_p\sigma \vdash t\theta \simeq t \quad (\text{dom}\theta \cap \text{fvar}(t) = \emptyset)$.

PROOF by easy reasoning in $\lambda_p\sigma$. \square

In the following we give an embedding $(\cdot)^{CL}$ of the system $\lambda_p\sigma$ into partial combinatory logic \mathbf{CL}_p . This embedding is made possible by a careful concept of substitution in the system $\lambda_p\sigma$ corresponding to substitution in \mathbf{CL}_p . As an immediate consequence of this interpretation we get that the recursion theoretic model *PRO* and the normal term model *CNT* are models of $\lambda_p\sigma$. This makes $\lambda_p\sigma$ into a system with a reasonable computational and constructive meaning.

Let us first define a \mathbf{CL}_p term t^{CL} for each $\lambda_p\sigma$ term t . The definition is by induction on the complexity of t .

Definition 20 (t^{CL})

1. If t is a variable, then $t^{CL} := t$.
2. If t is the term $(\lambda x.s)$, then $t^{CL} := (\lambda^*x.s^{CL})$.
3. If t is the term (t_1t_2) , then $t^{CL} := (t_1^{CL}t_2^{CL})$.
4. If t is the term $(s\theta)$ where $\theta = \{s_1/x_1, \dots, s_n/x_n\}$, then t^{CL} is the term $s^{CL}[s_1^{CL}/x_1, \dots, s_n^{CL}/x_n]$.

Once $(\cdot)^{CL}$ is defined for terms of $\lambda_p\sigma$, the translation for formulas is uniquely determined by the requirement that it commutes with $=$, \downarrow , the logical connectives, and the quantifiers. Hence, we have a \mathbf{CL}_p formula A^{CL} for each $\lambda_p\sigma$ formula A . From the definition of $(\cdot)^{CL}$ it is immediate that $\text{fvar}(t^{CL}) = \text{fvar}(t)$, $\text{fvar}(A^{CL}) = \text{fvar}(A)$, and $A\{t_1/x_1, \dots, t_n/x_n\}^{CL} = A^{CL}[t_1^{CL}/x_1, \dots, t_n^{CL}/x_n]$.

We are ready to state the embedding theorem.

Theorem 21 *We have for all $\lambda_p\sigma$ formulas A :*

$$\lambda_p\sigma \vdash A \implies \mathbf{CL}_p \vdash A^{CL}.^2$$

PROOF by induction on the length of a proof of A in $\lambda_p\sigma$. The propositional axioms do not cause any problems, of course. Also the quantifier axioms are easily handled using the properties of $(\cdot)^{CL}$ mentioned above. The equality axioms are trivial, and in order to establish the translation of $(\lambda x.t)\downarrow$ one makes immediate use of Lemma 8. The substitution axioms are easily verified, too. The extended β axiom (15) is treated in exactly the same way as in the proof of Lemma 11. Finally, it is trivial to check the inference rules. \square

In the sequel we show that $\lambda_p\sigma$ also includes \mathbf{CL}_p by giving an embedding $(\cdot)^\lambda$ from \mathbf{CL}_p into $\lambda_p\sigma$. We first give the translation $(\cdot)^\lambda$ for terms of \mathbf{CL}_p .

Definition 22 (t^λ)

1. If t is a variable, then $t^\lambda := t$.
2. If t is the constant \mathbf{k} , then $t^\lambda := \lambda uv.u$.
3. If t is the constant \mathbf{s} , then $t^\lambda := \lambda uvw.uw(vw)$.
4. If t is the term (t_1t_2) , then $t^\lambda := (t_1^\lambda t_2^\lambda)$.

For formulas, $(\cdot)^\lambda$ is given in the obvious way, i.e. $(\cdot)^\lambda$ commutes with $=$, \downarrow , the logical connectives, and the quantifiers. Again it is obvious that $fvar(t^\lambda) = fvar(t)$ and $fvar(A^\lambda) = fvar(A)$.

Lemma 23 *We have for all \mathbf{CL}_p terms t and s :*

$$\lambda_p\sigma \vdash t[s/x]^\lambda \simeq t^\lambda\{s^\lambda/x\}.$$

PROOF by straightforward induction on the complexity of t . Essential use is made of the substitution axioms, in particular axiom (13). \square

Corollary 24 *We have for all \mathbf{CL}_p formulas A , and for all \mathbf{CL}_p terms t :*

$$\lambda_p\sigma \vdash A[t/x]^\lambda \leftrightarrow A^\lambda\{t^\lambda/x\}.$$

PROOF by an easy induction on the complexity of A using the above lemma. \square

²Notice that the converse of this theorem does not hold. To see this, take e.g. for A the formula $x = y \rightarrow \lambda z.x = \lambda z.y$.

Lemma 25

1. $\lambda_p \sigma \vdash (\mathbf{k}xy \simeq x)^\lambda$.
2. $\lambda_p \sigma \vdash (\mathbf{s}xyz \simeq xz(yz))^\lambda$.
3. $\lambda_p \sigma \vdash (\mathbf{s}xy \downarrow)^\lambda$.

PROOF We only prove (1). The proof of (2) and (3) is similar. As we will see, essential use is made of the *extended* β axiom (15). First of all we have

$$(\lambda uv.u)x \simeq (\lambda v.u)\{x/u\}, \quad (1)$$

which by axiom (15) immediately implies

$$(\lambda uv.u)xy \simeq (\lambda v.u)\{x/u\}y \simeq u\{x/u, y/v\}. \quad (2)$$

Furthermore, we have

$$u\{x/u, y/v\} \simeq x. \quad (3)$$

This finishes the proof of our claim. \square

Lemma 26 *We have for all \mathbf{CL}_p formulas A :*

$$\mathbf{CL}_p \vdash A \implies \lambda_p \sigma \vdash A^\lambda.$$

PROOF by induction on the length of a proof of A in \mathbf{CL}_p . Again the propositional axioms are trivial. The translation of the quantifier axioms is provable in $\lambda_p \sigma$ by Corollary 24. The same corollary also helps in establishing the equality axiom (6). The strictness axioms (9) and the axioms for a partial combinatory algebra were already treated in Lemma 25. The inference rules of \mathbf{CL}_p readily translate into inference rules of $\lambda_p \sigma$. \square

The converse of the above lemma also holds.

Lemma 27 *We have for all \mathbf{CL}_p formulas A :*

$$\lambda_p \sigma \vdash A^\lambda \implies \mathbf{CL}_p \vdash A.$$

PROOF We define a modification $(\cdot)^\bullet$ of the translation $(\cdot)^{CL}$ from $\lambda_p \sigma$ into \mathbf{CL}_p . $(\cdot)^\bullet$ is defined in the same way as $(\cdot)^{CL}$, except that it uses the more complicated coding λ^\bullet of λ abstraction instead of λ^* . The term $\lambda^\bullet x.t$ is inductively defined as follows.

1. If t is the variable x , then $\lambda^\bullet x.t := \mathbf{s}kk$.

2. If t is a variable different from x or a constant, then $\lambda^{\bullet}x.t := \mathbf{k}t$.
3. If t is the term (sx) , where $s \in \{y, \mathbf{k}, \mathbf{s}, \mathbf{s}y\}$ and $y \neq x$, then $\lambda^{\bullet}x.t := s$.
4. If t is the term (t_1t_2) , and if (3) does not apply, then $\lambda^{\bullet}x.t := \mathbf{s}(\lambda^{\bullet}x.t_1)(\lambda^{\bullet}x.t_2)$.

One easily verifies that $(\mathbf{k}^{\lambda})^{\bullet} = \mathbf{k}$ and $(\mathbf{s}^{\lambda})^{\bullet} = \mathbf{s}$. As an immediate consequence we have $(t^{\lambda})^{\bullet} = t$ and $(A^{\lambda})^{\bullet} = A$ for all \mathbf{CL}_p terms t and all \mathbf{CL}_p formulas A , respectively. Furthermore, Lemma 8 holds for λ^{\bullet} , too. In particular, we have $\lambda^{\bullet}x.t \downarrow$ for all \mathbf{CL}_p terms t . We can, therefore, establish Theorem 21 for $(\cdot)^{\bullet}$ instead of $(\cdot)^{CL}$. Hence, we have

$$\lambda_p\sigma \vdash A \quad \Longrightarrow \quad \mathbf{CL}_p \vdash A^{\bullet}$$

for all $\lambda_p\sigma$ formulas A . Now the claim of the lemma immediately follows from the fact that $(\cdot)^{\bullet}$ is the inverse of $(\cdot)^{\lambda}$, as we have mentioned above. \square

Here is the final embedding theorem.

Theorem 28 *We have for all \mathbf{CL}_p formulas A :*

$$\mathbf{CL}_p \vdash A \quad \Longleftrightarrow \quad \lambda_p\sigma \vdash A^{\lambda}.$$

PROOF Immediate from the previous two lemmas. \square

4 Confluent reductions

Once we have introduced the system $\lambda_p\sigma$, it is natural to study the corresponding reduction relation on $\lambda_p\sigma$ terms. We now define a binary relation \triangleright on $\lambda_p\sigma$ terms, which reflects a directed equality relation for the system $\lambda_p\sigma$. \triangleright is defined inductively as follows.

Definition 29 The relation \triangleright between $\lambda_p\sigma$ terms is generated by the following clauses (1)–(13).

A. IDENTITY

$$(1) \quad t \triangleright t$$

B. β REDUCTIONS

$$(2) \quad (\lambda x.t)s \triangleright t\{s/x\}$$

$$(3) \quad (\lambda x.t)\theta s \triangleright t(s/x \cdot \theta)$$

C. SUBSTITUTION REDUCTIONS

- (4) $x\theta \triangleright t \quad (t/x \in \theta)$
- (5) $(ts)\theta \triangleright (t\theta)(s\theta)$
- (6) $(t\theta)\sigma \triangleright t(\theta\sigma)$
- (7) $t(s/x \cdot \theta) \triangleright t\theta \quad (x \notin \text{fvar}(t) \cup \text{dom}\theta)$
- (8) $t\varepsilon \triangleright t$

D. STRUCTURAL RULES

- (9) $t \triangleright s \implies r(t/x \cdot \theta) \triangleright r(s/x \cdot \theta)$
- (10) $t \triangleright s \implies t\theta \triangleright s\theta$
- (11) $t \triangleright s \implies rt \triangleright rs$
- (12) $t \triangleright s \implies tr \triangleright sr$

E. TRANSITIVITY

- (13) $t \triangleright s, s \triangleright r \implies t \triangleright r$

Notice that we have to state two clauses for β reduction, since (2) is no longer derivable from (3) in the context of reductions. Furthermore, it should be observed again that we do not have the rule (ξ),

$$(\xi) \frac{t \triangleright s}{\lambda x.t \triangleright \lambda x.s}$$

among the structural rules (D).

Remark 30 We want to stress that the reduction relation \triangleright does not take into consideration partiality in $\lambda_p\sigma$. Hence, \triangleright rather corresponds to a total version of $\lambda_p\sigma$. This is, however, in complete analogy to the system \mathbf{CL}_p , where truly partial term models are constructed using special reduction strategies of a *total* reduction relation (cf. p.7), and partiality is reflected by non-terminating reduction sequences. Summarizing, \triangleright provides a general term reduction framework giving rise to partial and total term models for $\lambda_p\sigma$.

In the following \triangleright_σ denotes the restriction of \triangleright to substitution reductions, i.e. $t \triangleright_\sigma s$ holds if and only if there is a derivation of $t \triangleright s$ according to the above clauses, which does not use (2) and (3). Analogously, \triangleright_β is the restriction of \triangleright to β reductions.

Finally, we write $t \triangleright^1 s$ if $t \triangleright s$ is derivable from (2)-(12), i.e. \triangleright^1 denotes one step reduction. The relations \triangleright_σ^1 and \triangleright_β^1 are defined in the same way.

In the sequel we want to show that \triangleright satisfies the Church Rosser property. As usual, this will guarantee that all terminating rewrite sequences yield identical results, i.e. we will have uniqueness of normal forms. All attempts to find a direct proof for the confluence of \triangleright failed. In particular, parallelization does not seem to work in order to show confluence of $\lambda_p\sigma$. Instead we will make use of an interpretation technique due to Hardin, which was identified in [12]. This method has subsequently been used several times in order to show confluence for systems of explicit substitutions.

A first step towards the proof of the Church Rosser property for \triangleright is to show that \triangleright_σ is Church Rosser. In order to apply an old result by Newman [25], we establish that \triangleright_σ^1 is weakly Church Rosser and wellfounded.

Lemma 31 \triangleright_σ^1 is weakly Church Rosser, i.e. we have for all terms t, t_1, t_2 : If $t \triangleright_\sigma^1 t_1$ and $t \triangleright_\sigma^1 t_2$, then there is a term t_3 such that $t_1 \triangleright_\sigma t_3$ and $t_2 \triangleright_\sigma t_3$.

PROOF One shows by a straightforward, but tedious induction on the length of a derivation of $t \triangleright_\sigma^1 t_1$ that for all $t \triangleright_\sigma^1 t_2$ there exists a t_3 such that $t_1 \triangleright_\sigma t_3$ and $t_2 \triangleright_\sigma t_3$. \square

In order to prove that \triangleright_σ^1 is wellfounded, we define a measure function Ψ from the $\lambda_p\sigma$ terms and $\lambda_p\sigma$ substitutions to ω .

Definition 32 $(\Psi(t); \Psi(\theta))$

1. If t is a variable, then $\Psi(t) := 1$.
2. If t is the term $(\lambda x.s)$, then $\Psi(t) := 1$.
3. If t is the term $(t_1 t_2)$, then $\Psi(t) := \Psi(t_1) + \Psi(t_2) + 1$.
4. If t is the term $(s\theta)$, then $\Psi(t) := \Psi(s) \cdot (\Psi(\theta) + 1)$.
5. If θ is the substitution $\{t_1/x_1, \dots, t_n/x_n\}$, then $\Psi(\theta) := \Psi(t_1) + \dots + \Psi(t_n) + 1$.

Notice that $\Psi(t) > 0$ for all terms t and $\Psi(\theta) > 0$ for all substitutions θ , e.g. we have $\Psi(\varepsilon) = 1$.

The composition $(\theta\sigma)$ of two substitutions θ and σ is bounded as follows w.r.t. Ψ .

Lemma 33 We have for all substitutions θ and σ :

$$\Psi(\theta\sigma) < \Psi(\theta) \cdot (\Psi(\sigma) + 1).$$

PROOF by an easy calculation. \square

We are ready to prove that Ψ is strictly decreasing on \triangleright_σ^1 .

Lemma 34 *We have for all terms t and s :*

$$t \triangleright_\sigma^1 s \implies \Psi(t) > \Psi(s).$$

PROOF by a straightforward induction on the length of a derivation of $t \triangleright_\sigma^1 s$. Let us only discuss the substitution reduction (6), i.e. $t = (r\theta)\sigma$ and $s = r(\theta\sigma)$ for some term r . Then we have

$$\Psi((r\theta)\sigma) = \Psi(r) \cdot (\Psi(\theta) + 1) \cdot (\Psi(\sigma) + 1) > \Psi(r) \cdot [\Psi(\theta) \cdot (\Psi(\sigma) + 1) + 1].$$

But $\Psi(r) \cdot [\Psi(\theta) \cdot (\Psi(\sigma) + 1) + 1] > \Psi(r) \cdot [\Psi(\theta\sigma) + 1]$ by the previous lemma. The claim is proved, since $\Psi(r) \cdot [\Psi(\theta\sigma) + 1] = \Psi(r(\theta\sigma))$. \square

Corollary 35 \triangleright_σ^1 *is wellfounded.*

Together we have established the following theorem.

Theorem 36 \triangleright_σ *is Church Rosser.*

PROOF The theorem is immediate from Lemma 31 and Corollary 35 and a result by Newman [25] saying that a reduction relation, which is weakly Church Rosser and wellfounded satisfies the full Church Rosser property. \square

Corollary 37 *Every $\lambda_p\sigma$ term has a unique substitution normal form.*

In the following we denote the substitution normal form of a term t with $\Sigma(t)$. A substitution $\theta = \{t_1/x_1, \dots, t_n/x_n\}$ is in substitution normal form, if for all i with $1 \leq i \leq n$ the term t_i is in substitution normal form. Analogously, $\Sigma(\theta)$ denotes the substitution normal form of a substitution θ .

As a further step towards the Church Rosser property of \triangleright we make use of a relation \triangleright_{β_n} , which corresponds to β reduction on terms in substitution normal form, i.e. terms satisfying $\Sigma(t) = t$. We define \triangleright_{β_n} via its “parallel” version \multimap_{β_n} , i.e. \triangleright_{β_n} will be the transitive closure of \multimap_{β_n} . Then the Church Rosser property of \multimap_{β_n} carries over to \triangleright_{β_n} by a well-known diagram chase.

For notational convenience we define \multimap_{β_n} on substitutions, too.

Definition 38 The relation \multimap_{β_n} between $\lambda_p\sigma$ terms and $\lambda_p\sigma$ substitutions in substitution normal form is simultaneously generated by the following clauses (1)–(6):

A. TERMS

- (1) $t \xrightarrow{\beta_n} t$
- (2) $s \xrightarrow{\beta_n} s' \implies (\lambda x.t)s \xrightarrow{\beta_n} \Sigma(t\{s'/x\})$
- (3) $s \xrightarrow{\beta_n} s', \theta \xrightarrow{\beta_n} \theta' \implies (\lambda x.t)\theta s \xrightarrow{\beta_n} \Sigma(t(s'/x \cdot \theta'))$
- (4) $\theta \xrightarrow{\beta_n} \theta' \implies (\lambda x.t)\theta \xrightarrow{\beta_n} (\lambda x.t)\theta'$
- (5) $t \xrightarrow{\beta_n} t', s \xrightarrow{\beta_n} s' \implies ts \xrightarrow{\beta_n} t's'$

B. SUBSTITUTIONS

- (6) $t_1 \xrightarrow{\beta_n} s_1, \dots, t_n \xrightarrow{\beta_n} s_n \implies \{t_1/x_1, \dots, t_n/x_n\} \xrightarrow{\beta_n} \{s_1/x_1, \dots, s_n/x_n\}$

Definition 39 Let \triangleright_{β_n} be the transitive closure of $\xrightarrow{\beta_n}$.

Lemma 40

1. If $t \xrightarrow{\beta_n} t'$ then we have $\Sigma(t\theta) \xrightarrow{\beta_n} \Sigma(t'\theta')$ for all $\theta \xrightarrow{\beta_n} \theta'$.
2. If $\theta \xrightarrow{\beta_n} \theta'$ then we have $\Sigma(\theta\sigma) \xrightarrow{\beta_n} \Sigma(\theta'\sigma')$ for all $\sigma \xrightarrow{\beta_n} \sigma'$.

PROOF (1) and (2) are proved simultaneously by induction on the complexity of t and θ , respectively. \square

The next lemma says that $\xrightarrow{\beta_n}$ is Church Rosser on terms and substitutions.

Lemma 41

1. If $t \xrightarrow{\beta_n} t_1$ then for all $t \xrightarrow{\beta_n} t_2$ there is a t_3 such that $t_1 \xrightarrow{\beta_n} t_3$ and $t_2 \xrightarrow{\beta_n} t_3$.
2. If $\theta \xrightarrow{\beta_n} \theta_1$ then for all $\theta \xrightarrow{\beta_n} \theta_2$ there is a θ_3 such that $\theta_1 \xrightarrow{\beta_n} \theta_3$ and $\theta_2 \xrightarrow{\beta_n} \theta_3$.

PROOF We prove (1) and (2) simultaneously by induction on $t \xrightarrow{\beta_n} t_1$ and $\theta \xrightarrow{\beta_n} \theta_1$, respectively. Let us first prove (1). According to $t \xrightarrow{\beta_n} t_1$ we can distinguish the following five cases:

- (1) $t \xrightarrow{\beta_n} t_1$ is $t \xrightarrow{\beta_n} t$. Then we can choose $t_3 := t_2$.
- (2) $t \xrightarrow{\beta_n} t_1$ is $(\lambda x.r)s \xrightarrow{\beta_n} \Sigma(r\{s'/x\})$ and is a consequence of $s \xrightarrow{\beta_n} s'$. According to $t \xrightarrow{\beta_n} t_2$ we can distinguish the following two subcases:
 - (2.1) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r)s \xrightarrow{\beta_n} (\lambda x.r)s''$ and is a consequence of $s \xrightarrow{\beta_n} s''$. By the induction hypothesis there is a term s''' with $s' \xrightarrow{\beta_n} s'''$ and $s'' \xrightarrow{\beta_n} s'''$. If we take

$t_3 := \Sigma(r\{s'''/x\})$ then obviously $t_2 \xrightarrow{\beta_n} t_3$. By the previous lemma we also have $t_1 \xrightarrow{\beta_n} t_3$.

(2.2) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r)s \xrightarrow{\beta_n} \Sigma(r\{s''/x\})$ and is a consequence of $s \xrightarrow{\beta_n} s''$. By the induction hypothesis there exists a term s''' with $s' \xrightarrow{\beta_n} s'''$ and $s'' \xrightarrow{\beta_n} s'''$. Let t_3 be $\Sigma(r\{s'''/x\})$. Using the previous lemma one easily verifies $t_1 \xrightarrow{\beta_n} t_3$ and $t_2 \xrightarrow{\beta_n} t_3$.

(3) $t \xrightarrow{\beta_n} t_1$ is $(\lambda x.r)\theta s \xrightarrow{\beta_n} \Sigma(r\{s'/x \cdot \theta'\})$ and is a consequence of $s \xrightarrow{\beta_n} s'$ and $\theta \xrightarrow{\beta_n} \theta'$. According to $t \xrightarrow{\beta_n} t_2$ we can distinguish the following two subcases:

(3.1) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r)\theta s \xrightarrow{\beta_n} (\lambda x.r)\theta'' s''$ and is a consequence of $\theta \xrightarrow{\beta_n} \theta''$ and $s \xrightarrow{\beta_n} s''$. By the induction hypothesis there are s''' and θ''' with $s' \xrightarrow{\beta_n} s'''$, $s'' \xrightarrow{\beta_n} s'''$ and $\theta' \xrightarrow{\beta_n} \theta'''$, $\theta'' \xrightarrow{\beta_n} \theta'''$. If we take $t_3 := \Sigma(r\{s'''/x \cdot \theta'''\})$ then obviously $t_2 \xrightarrow{\beta_n} t_3$. By the fact that $s'/x \cdot \theta' \xrightarrow{\beta_n} s'''/x \cdot \theta'''$ and the previous lemma we also have $t_1 \xrightarrow{\beta_n} t_3$.

(3.2) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r)\theta s \xrightarrow{\beta_n} \Sigma(r\{s''/x \cdot \theta''\})$ and is a consequence of $s \xrightarrow{\beta_n} s''$ and $\theta \xrightarrow{\beta_n} \theta''$. By the induction hypothesis there are s''' and θ''' with $s' \xrightarrow{\beta_n} s'''$, $s'' \xrightarrow{\beta_n} s'''$ and $\theta' \xrightarrow{\beta_n} \theta'''$, $\theta'' \xrightarrow{\beta_n} \theta'''$. Let $t_3 := \Sigma(r\{s'''/x \cdot \theta'''\})$. Then, by the previous lemma, we have $t_1 \xrightarrow{\beta_n} t_3$ and $t_2 \xrightarrow{\beta_n} t_3$, since $s'/x \cdot \theta' \xrightarrow{\beta_n} s'''/x \cdot \theta'''$ and $s''/x \cdot \theta'' \xrightarrow{\beta_n} s'''/x \cdot \theta'''$.

(4) $t \xrightarrow{\beta_n} t_1$ is $(\lambda x.r)\theta \xrightarrow{\beta_n} (\lambda x.r)\theta'$ and is a consequence of $\theta \xrightarrow{\beta_n} \theta'$. Then $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r)\theta \xrightarrow{\beta_n} (\lambda x.r)\theta''$ and is a consequence of $\theta \xrightarrow{\beta_n} \theta''$. By the induction hypothesis there is a θ''' with $\theta' \xrightarrow{\beta_n} \theta'''$ and $\theta'' \xrightarrow{\beta_n} \theta'''$. The claim holds for $t_3 := (\lambda x.r)\theta'''$.

(5) $t \xrightarrow{\beta_n} t_1$ is $rs \xrightarrow{\beta_n} r's'$ and is a consequence of $r \xrightarrow{\beta_n} r'$ and $s \xrightarrow{\beta_n} s'$. According to $t \xrightarrow{\beta_n} t_2$ we can distinguish the following three subcases:

(5.1) $t \xrightarrow{\beta_n} t_2$ is $rs \xrightarrow{\beta_n} r''s''$ and is a consequence of $r \xrightarrow{\beta_n} r''$ and $s \xrightarrow{\beta_n} s''$. By the induction hypothesis there are terms r''' and s''' with $r' \xrightarrow{\beta_n} r'''$, $r'' \xrightarrow{\beta_n} r'''$ and $s' \xrightarrow{\beta_n} s'''$, $s'' \xrightarrow{\beta_n} s'''$. The claim immediately follows for $t_3 := r'''s'''$.

(5.2) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r'')s \xrightarrow{\beta_n} \Sigma(r''\{s''/x\})$ and is a consequence of $s \xrightarrow{\beta_n} s''$. By the induction hypothesis there is an s''' with $s' \xrightarrow{\beta_n} s'''$ and $s'' \xrightarrow{\beta_n} s'''$. Furthermore, it is clear that $r = r' = (\lambda x.r'')$. If we take $t_3 := \Sigma(r''\{s'''/x\})$ then obviously $t_1 \xrightarrow{\beta_n} t_3$. By the previous lemma we also have $t_2 \xrightarrow{\beta_n} t_3$.

(5.3) $t \xrightarrow{\beta_n} t_2$ is $(\lambda x.r'')\theta s \xrightarrow{\beta_n} \Sigma(r''\{s''/x \cdot \theta''\})$ and is a consequence of $s \xrightarrow{\beta_n} s''$ and $\theta \xrightarrow{\beta_n} \theta''$. By the induction hypothesis there is an s''' with $s' \xrightarrow{\beta_n} s'''$ and $s'' \xrightarrow{\beta_n} s'''$. Furthermore, $r \xrightarrow{\beta_n} r'$ has the form $(\lambda x.r'')\theta \xrightarrow{\beta_n} (\lambda x.r'')\theta'$ and is a consequence of $\theta \xrightarrow{\beta_n} \theta'$. By the induction hypothesis there is a θ''' with $\theta' \xrightarrow{\beta_n} \theta'''$ and $\theta'' \xrightarrow{\beta_n} \theta'''$. If we take $t_3 := \Sigma(r''\{s'''/x \cdot \theta'''\})$ then obviously $t_1 \xrightarrow{\beta_n} t_3$. By the previous lemma we also have $t_2 \xrightarrow{\beta_n} t_3$.

This finishes the proof of (1). The proof of (2) is straightforward. Assume $\theta \xrightarrow{\beta_n} \theta_1$ and $\theta \xrightarrow{\beta_n} \theta_2$, where

$$\begin{aligned}\theta &= \{t_1/x_1, \dots, t_n/x_n\}, \\ \theta_1 &= \{s_1/x_1, \dots, s_n/x_n\}, \\ \theta_2 &= \{s'_1/x_1, \dots, s'_n/x_n\}\end{aligned}$$

and $t_i \xrightarrow{\beta_n} s_i$, $t_i \xrightarrow{\beta_n} s'_i$ for $1 \leq i \leq n$. By the induction hypothesis there are terms r_i with $s_i \xrightarrow{\beta_n} r_i$ and $s'_i \xrightarrow{\beta_n} r_i$ for $1 \leq i \leq n$. If we take $\theta_3 := \{r_1/x_1, \dots, r_n/x_n\}$, then we immediately get $\theta_1 \xrightarrow{\beta_n} \theta_3$ and $\theta_2 \xrightarrow{\beta_n} \theta_3$. \square

The proof of the Church Rosser property of \triangleright_{β_n} is complete.

Theorem 42 \triangleright_{β_n} is Church Rosser.

PROOF By the previous lemma, $\xrightarrow{\beta_n}$ is Church Rosser. By Definition 39, \triangleright_{β_n} is the transitive closure of $\xrightarrow{\beta_n}$, which, by a simple diagram chase, implies that \triangleright_{β_n} is confluent, too. \square

The last step towards the Church Rosser property of \triangleright is to show that

$$t \triangleright_{\beta}^1 s \implies \Sigma(t) \triangleright_{\beta_n} \Sigma(s)$$

holds for all $\lambda_p\sigma$ terms t and s . Then the confluence of \triangleright will be an immediate consequence. In order to establish the above claim, we have to define another intermediate relation, the reduction relation $\xrightarrow{\beta}$.

Definition 43 The relation $\xrightarrow{\beta}$ between $\lambda_p\sigma$ terms and $\lambda_p\sigma$ substitutions is given by the following clauses (1)–(8):

A. TERMS

- (1) $t \xrightarrow{\beta} t$
- (2) $(\lambda x.t)s \xrightarrow{\beta} t\{s/x\}$
- (3) $(\lambda x.t)\theta s \xrightarrow{\beta} t(s/x \cdot \theta)$
- (4) $\theta \xrightarrow{\beta} \theta' \implies r\theta \xrightarrow{\beta} r\theta'$
- (5) $t \xrightarrow{\beta} s \implies t\theta \xrightarrow{\beta} s\theta$
- (6) $t \xrightarrow{\beta} s \implies rt \xrightarrow{\beta} rs$
- (7) $t \xrightarrow{\beta} s \implies tr \xrightarrow{\beta} sr$

B. SUBSTITUTIONS

$$(8) \quad t_1 \xrightarrow{\beta} s_1, \dots, t_n \xrightarrow{\beta} s_n \quad \Longrightarrow \quad \{t_1/x_1, \dots, t_n/x_n\} \xrightarrow{\beta} \{s_1/x_1, \dots, s_n/x_n\}$$

The relation $\xrightarrow{\beta}$ can be considered as an extended form of one step β reduction, where substitutions can be reduced in parallel.

Lemma 44 *We have for all terms t and s :*

$$t \xrightarrow{\beta} s \quad \Longrightarrow \quad \Sigma(t) \triangleright_{\beta_n} \Sigma(s).$$

PROOF We prove the claim by main induction on $\Psi(t)$ and side induction on the complexity of t . According to the structure of t , we can distinguish the following four cases:

(1) t is a variable. Then the claim is trivial.

(2) t is the term $(\lambda x.t')$. The claim is trivial, too.

(3) t is the term $(t_1 t_2)$. According to $t \xrightarrow{\beta} s$ we can distinguish the following four subcases:

(3.1) $t \xrightarrow{\beta} s$ is $t_1 t_2 \xrightarrow{\beta} t'_1 t_2$ and is a consequence of $t_1 \xrightarrow{\beta} t'_1$. By the side induction hypothesis we have $\Sigma(t_1) \triangleright_{\beta_n} \Sigma(t'_1)$. This implies

$$\Sigma(t_1 t_2) = \Sigma(t_1) \Sigma(t_2) \triangleright_{\beta_n} \Sigma(t'_1) \Sigma(t_2) = \Sigma(t'_1 t_2).$$

(3.2) $t \xrightarrow{\beta} s$ is $t_1 t_2 \xrightarrow{\beta} t_1 t'_2$ and is a consequence of $t_2 \xrightarrow{\beta} t'_2$. This case is treated in the same way as (3.1).

(3.3) $t \xrightarrow{\beta} s$ is $(\lambda x.r)r' \xrightarrow{\beta} r\{r'/x\}$. Then we have

$$\Sigma((\lambda x.r)r') = \Sigma(\lambda x.r) \Sigma(r') = (\lambda x.r) \Sigma(r') \triangleright_{\beta_n} \Sigma(r\{\Sigma(r')/x\}) = \Sigma(r\{r'/x\}).$$

(3.4) $t \xrightarrow{\beta} s$ is $(\lambda x.r)\theta r' \xrightarrow{\beta} r(r'/x \cdot \theta)$. First assume that $fvar(\lambda x.r) \cap dom \theta \neq \emptyset$. Then we have for $\theta' := \theta \upharpoonright (\lambda x.r)$

$$\Sigma((\lambda x.r)\theta r') = (\lambda x.r) \Sigma(\theta') \Sigma(r') \triangleright_{\beta_n} \Sigma(r(\Sigma(r')/x \cdot \Sigma(\theta'))) = \Sigma(r(r'/x \cdot \theta)).$$

The case $fvar(\lambda x.r) \cap dom \theta = \emptyset$ is treated in a similar way.

(4) t is the term $(t'\theta)$. According to the structure of t' we can distinguish the following four subcases:

(4.1) t' is the variable x . Then s is of the form $x\theta'$, where $\theta \xrightarrow{\beta} \theta'$. If $x \notin dom \theta$ then $\Sigma(x\theta) = x = \Sigma(x\theta')$ and there is nothing to prove. Therefore, assume $r/x \in \theta$,

$r'/x \in \theta'$ and $r \xrightarrow{\beta} r'$ for some terms r and r' . Then it is $x\theta \triangleright_{\sigma}^1 r$ and by the main induction hypothesis we have

$$\Sigma(x\theta) = \Sigma(r) \triangleright_{\beta_n} \Sigma(r') = \Sigma(x\theta').$$

(4.2) t' is the term $(\lambda x.t'')$. Then s is of the form $(\lambda x.t'')\theta'$, where $\theta \xrightarrow{\beta} \theta'$. If $fvar(\lambda x.t'') \cap dom\theta = \emptyset$ then $\Sigma((\lambda x.t'')\theta) = \lambda x.t'' = \Sigma((\lambda x.t'')\theta')$ and there is nothing to prove. Otherwise, by the side induction hypothesis, we have

$$\Sigma((\lambda x.t'')\theta) = (\lambda x.t'')(\theta \upharpoonright (\lambda x.t'')) \triangleright_{\beta_n} (\lambda x.t'')(\theta' \upharpoonright (\lambda x.t'')) = \Sigma((\lambda x.t'')\theta').$$

(4.3) t' is the term $(t'_1 t'_2)$. According to $t'\theta \xrightarrow{\beta} s$, we can distinguish the following five subcases:

(4.3.1) $t'\theta \xrightarrow{\beta} s$ is $(t'_1 t'_2)\theta \xrightarrow{\beta} (t''_1 t''_2)\theta$ and is a consequence of $t'_1 \xrightarrow{\beta} t''_1$. Then it is $(t'_1 t'_2)\theta \triangleright_{\sigma}^1 (t'_1\theta)(t'_2\theta)$ and by the main induction hypothesis we have

$$\Sigma((t'_1 t'_2)\theta) = \Sigma((t'_1\theta)(t'_2\theta)) \triangleright_{\beta_n} \Sigma((t''_1\theta)(t''_2\theta)) = \Sigma((t''_1 t''_2)\theta).$$

(4.3.2) $t'\theta \xrightarrow{\beta} s$ is $(t'_1 t'_2)\theta \xrightarrow{\beta} (t'_1 t''_2)\theta$ and is a consequence of $t'_2 \xrightarrow{\beta} t''_2$. This case is treated in the same way as (4.3.1).

(4.3.3) $t'\theta \xrightarrow{\beta} s$ is $(t'_1 t'_2)\theta \xrightarrow{\beta} (t'_1 t'_2)\theta'$ and is a consequence of $\theta \xrightarrow{\beta} \theta'$. Then it is $(t'_1 t'_2)\theta \triangleright_{\sigma}^1 (t'_1\theta)(t'_2\theta)$ and by the main induction hypothesis we get $\Sigma(t'_1\theta) \triangleright_{\beta_n} \Sigma(t'_1\theta')$ and $\Sigma(t'_2\theta) \triangleright_{\beta_n} \Sigma(t'_2\theta')$. This implies

$$\Sigma((t'_1 t'_2)\theta) = \Sigma((t'_1\theta)(t'_2\theta)) = \Sigma(t'_1\theta)\Sigma(t'_2\theta) \triangleright_{\beta_n} \Sigma(t'_1\theta')\Sigma(t'_2\theta') = \Sigma((t'_1 t'_2)\theta').$$

(4.3.4) $t'\theta \xrightarrow{\beta} s$ is $((\lambda x.r)r')\theta \xrightarrow{\beta} r\{r'/x\}\theta$. First assume that $fvar(\lambda x.r) \cap dom\theta \neq \emptyset$. Then we have for $\theta' := \theta \upharpoonright (\lambda x.r)$

$$\Sigma(((\lambda x.r)r')\theta) = (\lambda x.r)\Sigma(\theta')\Sigma(r'\theta) \triangleright_{\beta_n} \Sigma(r(\Sigma(r'\theta)/x \cdot \Sigma(\theta'))) = \Sigma(r\{r'/x\}\theta).$$

The case $fvar(\lambda x.r) \cap dom\theta = \emptyset$ is treated in a similar way.

(4.3.5) $t'\theta \xrightarrow{\beta} s$ is $((\lambda x.r)\sigma r')\theta \xrightarrow{\beta} r(r'/x \cdot \sigma)\theta$. Let us first assume that we have $fvar(\lambda x.r) \cap dom(\sigma\theta) \neq \emptyset$. Then we get for $\rho := (\sigma\theta) \upharpoonright (\lambda x.r)$

$$\Sigma(((\lambda x.r)\sigma r')\theta) = (\lambda x.r)\Sigma(\rho)\Sigma(r'\theta) \triangleright_{\beta_n} \Sigma(r(\Sigma(r'\theta)/x \cdot \Sigma(\rho))) = \Sigma(r(r'/x \cdot \sigma)\theta).$$

The case $fvar(\lambda x.r) \cap dom(\sigma\theta) = \emptyset$ is treated in a similar way.

(4.4) t' is the term $(t''\sigma)$. Since $t''\sigma\theta \triangleright_{\sigma}^1 t''(\sigma\theta)$ we can apply the main induction hypothesis to $t''(\sigma\theta)$. According to $t'\theta \xrightarrow{\beta} s$ we can distinguish the following three subcases:

(4.4.1) $t'\theta \xrightarrow{\beta} s$ is $t''\sigma\theta \xrightarrow{\beta} t''\sigma\theta'$ and is a consequence of $\theta \xrightarrow{\beta} \theta'$. Then, by the main induction hypothesis, we have

$$\Sigma(t''\sigma\theta) = \Sigma(t''(\sigma\theta)) \triangleright_{\beta_n} \Sigma(t''(\sigma\theta')) = \Sigma(t''\sigma\theta').$$

(4.4.2) $t'\theta \xrightarrow{\beta} s$ is $t''\sigma\theta \xrightarrow{\beta} t''\sigma'\theta$ and is a consequence of $\sigma \xrightarrow{\beta} \sigma'$. This case is treated in a very similar way as (4.4.1).

(4.4.3) $t'\theta \xrightarrow{\beta} s$ is $t''\sigma\theta \xrightarrow{\beta} t'''\sigma\theta$ and is a consequence of $t'' \xrightarrow{\beta} t'''$. Then, by the main induction hypothesis, we have

$$\Sigma(t''\sigma\theta) = \Sigma(t''(\sigma\theta)) \triangleright_{\beta_n} \Sigma(t'''(\sigma\theta)) = \Sigma(t'''\sigma\theta).$$

This finishes the proof of our claim. \square

Since $t \triangleright_{\beta}^1 s$ trivially implies $t \xrightarrow{\beta} s$, we get the following corollary.

Corollary 45 *We have for all terms t and s :*

$$t \triangleright_{\beta}^1 s \implies \Sigma(t) \triangleright_{\beta_n} \Sigma(s).$$

Corollary 46 *We have for all terms t and s :*

$$t \triangleright s \implies \Sigma(t) \triangleright_{\beta_n} \Sigma(s).$$

PROOF Assume $t \triangleright s$. Then there is a sequence of terms t_1, \dots, t_n so that we have $t_1 = t, t_n = s$ and for all $1 \leq i < n$

$$t_i \triangleright_{\sigma}^1 t_{i+1} \quad \text{or} \quad t_i \triangleright_{\beta}^1 t_{i+1}.$$

In the first case we have $\Sigma(t_i) = \Sigma(t_{i+1})$ and in the latter, by the previous corollary, $\Sigma(t_i) \triangleright_{\beta_n} \Sigma(t_{i+1})$. Together we immediately get $\Sigma(t) \triangleright_{\beta_n} \Sigma(s)$, since \triangleright_{β_n} is transitively closed. \square

We are ready to prove the confluence of \triangleright .

Theorem 47 \triangleright *is Church Rosser.*

PROOF Let t, t_1 and t_2 be $\lambda_p\sigma$ terms and assume that

$$t \triangleright t_1 \quad \text{and} \quad t \triangleright t_2. \tag{1}$$

Then the previous corollary immediately implies

$$\Sigma(t) \triangleright_{\beta_n} \Sigma(t_1) \quad \text{and} \quad \Sigma(t) \triangleright_{\beta_n} \Sigma(t_2). \tag{2}$$

By Theorem 42 we know that \triangleright_{β_n} is confluent, hence there is a $\lambda_p\sigma$ term s in substitution normal form satisfying

$$\Sigma(t_1) \triangleright_{\beta_n} s \quad \text{and} \quad \Sigma(t_2) \triangleright_{\beta_n} s. \quad (3)$$

Since $\triangleright_{\beta_n} \subset \triangleright$, we get from (3)

$$\Sigma(t_1) \triangleright s \quad \text{and} \quad \Sigma(t_2) \triangleright s, \quad (4)$$

which immediately implies

$$t_1 \triangleright \Sigma(t_1) \triangleright s \quad \text{and} \quad t_2 \triangleright \Sigma(t_2) \triangleright s. \quad (5)$$

The claim is established. \square

5 Conclusion

We have addressed some defects of the partial λ calculus λ_p as a constructive framework for partial functions. The drawbacks of λ_p become even more perspicuous in the light of Pezzoli's theorem (cf. Theorem 9). We have proposed a modification $\lambda_p\sigma$ of λ_p by explicit substitutions. The system $\lambda_p\sigma$ is embeddable into partial combinatory logic \mathbf{CL}_p , and therefore, inherits all its models. In particular, $\lambda_p\sigma$ has a standard interpretation in terms of ordinary recursion theory. We have studied a reduction relation for $\lambda_p\sigma$ and we have established a confluence result. The reduction relation gives rise to direct constructions of term models for $\lambda_p\sigma$. The detailed constructions will be discussed later.

As already mentioned, the theory of explicit substitutions has been treated before, primarily in connection with the implementation of functional programming languages. The main reference on weak calculi of explicit substitutions is [5]. In contrast to our approach, not only equality between terms, but also equality between substitutions has been axiomatized in most of the previous work on explicit substitutions (an exception is [20]). Although this can easily be achieved, the systems have much more axioms, and we think that – especially from a foundational point of view – the real concern is to axiomatize and control the notion of a substitution applied to a term, whereas equality between substitutions can be treated in the metalanguage. Furthermore, the previous systems of explicit substitutions are mainly term rewriting systems, and – this is the main difference to our $\lambda_p\sigma$ calculus – application in those systems is always total. The very purpose of our work, however, was to study questions of substitution in the context of partiality, and to design a more perspicuous version of the partial λ calculus, which has natural partial models and is equivalent to partial combinatory logic.

The question arises, why one should use a fairly complicated system like $\lambda_p\sigma$ at all, instead of the formally more simple partial combinatory logic \mathbf{CL}_p . We think that $\lambda_p\sigma$ has advantages over \mathbf{CL}_p . The main reason is that in the system \mathbf{CL}_p , the intuitive clarity of the λ notation is completely lost. Additionally, many mistakes in the literature concerning substitution in \mathbf{CL}_p suggest that it is also worth having an explicit treatment of substitution as in $\lambda_p\sigma$. We, therefore, think that $\lambda_p\sigma$ can serve as an adequate applicative basis for systems of explicit mathematics.

As already mentioned in the introduction, Stärk [27] has recently given a very natural relationship between the programming language SCHEME and the $\lambda_p\sigma$ calculus. In his approach, partiality of $\lambda_p\sigma$ is essential. The results of Stärk give further evidence for the foundational significance of the $\lambda_p\sigma$ calculus.

Acknowledgments

I am indebted to R. Kahle, who pointed out to me that the principle $(\star\star)$ is derivable in λ_p though it obviously does not hold in the recursion-theoretic model, and to E. Pezzoli for providing the crucial Theorem 9. I also thank A. S. Troelstra for calling my attention to explicit substitutions as well as S. Feferman, T. Hardin, G. Jäger, R. F. Stärk, and W. Th. Wolff for helpful comments on an earlier version of this paper.

References

- [1] ABADI, M., CARDELLI, L., CURIEN, P.-L., AND LEVY, J.-J. Explicit substitutions. *Journal of Functional Programming* 1, 4 (1991), 375–416.
- [2] BEESON, M. J. *Foundations of Constructive Mathematics: Metamathematical Studies*. Springer, Berlin, 1984.
- [3] BEESON, M. J. Proving programs and programming proofs. In *Logic, Methodology and Philosophy of Science VII*, Barcan Marcus et. al., Ed. North Holland, Amsterdam, 1986, pp. 51–82.
- [4] CURIEN, P.-L. An abstract framework for environment machines. *Theoretical Computer Science* 82 (1991), 389–402.
- [5] CURIEN, P.-L., HARDIN, T., AND LEVY, J.-J. Confluence properties of weak and strong calculi of explicit substitutions. Tech. Rep. 92-01, CEDRIC, 292 Rue Saint-Martin, 75141 Paris Cédex 03, France, 1992. Submitted to J.A.C.M.

- [6] FEFERMAN, S. A language and axioms for explicit mathematics. In *Algebra and Logic*, J. N. Crossley, Ed., vol. 450 of *Lecture Notes in Mathematics*. Springer, Berlin, 1975, pp. 87–139.
- [7] FEFERMAN, S. Constructive theories of functions and classes. In *Logic Colloquium '78*, M. Boffa, D. van Dalen, and K. McAloon, Eds. North Holland, Amsterdam, 1979, pp. 159–224.
- [8] FEFERMAN, S. Polymorphic typed lambda-calculi in a type-free axiomatic framework. In *Logic and Computation*, W. Sieg, Ed., vol. 106 of *Contemporary Mathematics*. American Mathematical Society, Providence, Rhode Island, 1990, pp. 101–136.
- [9] FEFERMAN, S. Logics for termination and correctness of functional programs. In *Logic from Computer Science*, Y. N. Moschovakis, Ed., vol. 21 of *MSRI Publications*. Springer, Berlin, 1991, pp. 95–127.
- [10] FEFERMAN, S. Logics for termination and correctness of functional programs II: Logics of strength PRA. In *Proof Theory*, P. Aczel, H. Simmons, and S. S. Wainer, Eds. Cambridge University Press, Cambridge, 1992, pp. 195–225.
- [11] FEFERMAN, S., AND JÄGER, G. Systems of explicit mathematics with non-constructive μ -operator. Part I. *Annals of Pure and Applied Logic* 65, 3 (1993), 243–263.
- [12] HARDIN, T. Confluence results for the pure strong categorical logic CCL: λ -calculi as subsystems of CCL. *Theoretical Computer Science* 65 (1989), 291–342.
- [13] HARDIN, T., AND LEVY, J.-J. A confluent calculus of substitutions. Tech. Rep. 90-11, CEDRIC, 292 Rue Saint-Martin, 75141 Paris Cédex 03, France, 1990.
- [14] HINDLEY, J. R. Combinatory reductions and lambda reductions compared. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik* 23 (1977), 169–180.
- [15] HINDLEY, J. R., AND LONGO, G. Lambda-calculus models and extensionality. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik* 26 (1980), 289–310.
- [16] JÄGER, G. Induction in the elementary theory of types and names. In *Computer Science Logic '87*, E. Börger et al., Ed., vol. 329 of *Lecture Notes in Computer Science*. Springer, Berlin, 1988, pp. 118–128.

- [17] JÄGER, G. Type theory and explicit mathematics. In *Logic Colloquium '87*, H.-D. Ebbinghaus, J. Fernandez-Prida, M. Garrido, M. Lascar, and M. R. Artalejo, Eds. North Holland, Amsterdam, 1989, pp. 117–135.
- [18] JÄGER, G., AND STRAHM, T. Totality in applicative theories. *Annals of Pure and Applied Logic*. To appear.
- [19] KLEENE, S. C. *Introduction to Metamathematics*. North Holland, Amsterdam, 1952.
- [20] LESCANNE, P., AND ROUYER-DEGLI, J. The calculus of explicit substitution λv . Tech. rep., CNRS and INRIA-Lorraine, Campus Scientifique, BP 239, 54506 Vandœuvre-lès-Nancy, France, March 1994.
- [21] MARTIN-LÖF, P. Substitution calculus. Unpublished notes, September 1992.
- [22] MARZETTA, M. A theory of types and names. Contributed paper, Logic Colloquium, Padova 23-30 August 1988.
- [23] MARZETTA, M. Universes in the theory of types and names. In *Computer Science Logic '92*, E. Börger et al., Ed., vol. 702 of *Lecture Notes in Computer Science*. Springer, Berlin, 1993, pp. 340–351.
- [24] MOGGI, E. *The Partial Lambda Calculus*. PhD thesis, University of Edinburgh, 1988.
- [25] NEWMAN, M. H. A. On theories with a combinatorial definition of “equivalence”. *Annals of Mathematics* 43, 2 (1942), 223–243.
- [26] PEZZOLI, E. Private communication, November 1993.
- [27] STÄRK, R. F. Applicative theories and SCHEME. In preparation, May 1994.
- [28] TASISTRO, A. Formulation of Martin-Löf’s theory of types with explicit substitutions. Tech. rep., Department of Computer Sciences, University of Göteborg, May 1993.
- [29] TROELSTRA, A. S., AND VAN DALEN, D. *Constructivism in Mathematics*, vol. II. North Holland, Amsterdam, 1988.
- [30] TROELSTRA, A. S., AND VAN DALEN, D. *Constructivism in Mathematics*, vol. I. North Holland, Amsterdam, 1988.