

# The Modal $\mu$ -Calculus Hierarchy over Restricted Classes of Transition Systems

Luca Alberucci<sup>1\*</sup> and Alessandro Facchini<sup>2,3\*\*</sup>

<sup>1</sup> University of Berne, Switzerland

<sup>2</sup> University of Lausanne, Switzerland

<sup>3</sup> University of Bordeaux 1, France

albe@iam.unibe.ch, alessandro.facchini@unil.ch

**Abstract.** We study the strictness of the modal  $\mu$ -calculus hierarchy over some restricted classes of transition systems. First, we show that the hierarchy is strict over reflexive frames. By proving the finite model theorem for reflexive systems the same results holds for finite models. Second, we prove that over transitive systems the hierarchy collapses to the alternation-free fragment. In order to do this the finite model theorem for transitive transition systems is also proved. Further, we verify that if symmetry is added to transitivity the hierarchy collapses to the purely modal fragment.

## 1 Introduction

The modal  $\mu$ -calculus is an extension of modal logics, with least and greatest fixpoint operators. The term “ $\mu$ -calculus” and the idea of extending modal logic with fixpoints appeared for the first time in the paper of Scott and De Bakker [SB69] and was further developed by others. Nowadays, the term “modal  $\mu$ -calculus” stands for the formal system introduced by Kozen [Koz83]. It is a powerful logic of programs subsuming dynamic and temporal logics like PDL, PLTL, CTL and CTL\*. Hence, it provides us with the capability of expressing and reasoning about assertions concerning “temporal” properties of dynamic (reactive and parallel) systems with potentially infinite behavior. We refer to Bradfield and Stirling’s tutorial article [BS01] or Stirling’s book [Sti01] for a thorough introduction to this system.

The standard semantics of the modal  $\mu$ -calculus is given by transition systems. As usual, formulae are interpreted as subsets of a system, the set of states where the property expressed by the formula holds. Many natural properties such as “there is an infinite path” can be expressed by a modal  $\mu$ -formula. Further, most such properties are given by formulae with alternation depth two, where the alternation depth is the number of non-trivial nestings of least and greatest

---

\* Research supported by the Hasler Foundation

\*\* Research supported by a grant from the Swiss National Science Foundation, n. 200021-116508: Project “Topological Complexity, Games, Logic and Automata”.

fixpoints. Nevertheless, it is mathematically interesting to see whether the expressive power of the modal  $\mu$ -calculus increases with the alternation depth. If this is the case then we have a strict hierarchy otherwise we have a collapse at some point.

By a result of Bradfield [Bra98<sup>1</sup>,Bra98<sup>2</sup>] the hierarchy over arbitrary transition systems is strict, a result independently proved by Lenzi in [Len96] but only for the positive fragment. Subsequently, Arnold showed in [Arn99] that the hierarchy is also strict over binary trees by using the Banach-Cacciopoli fixpoint theorem. His proof has been naturally extended to the class of all models by one of the authors in [Alb02].

Having seen the strictness over arbitrary transition systems, it can be asked whether the alternation hierarchy remains strict for restricted classes of transition systems such as those that are reflexive or those that are transitive. In the case of transitive systems, to our knowledge, the first attempt to answer this question is presented by Lenzi in [Len06]. There, he shows that on transitive frames every Büchi automaton is equivalent to a co-Büchi automaton, and conversely<sup>4</sup>. This implies that over transitive frames the modal  $\mu$ -calculus collapses to the level of Büchi automata (and to co-Büchi automata). Because, for example, well-foundedness is not definable in the modal fragment, the hierarchy is non-trivial. Thus, since over arbitrary graphs the intersection of Büchi and co-Büchi automata corresponds to the alternation-free fragment, Lenzi conjectured that the full modal  $\mu$ -calculus collapses to the alternation-free fragment ([Len08]). It is interesting to note that Visser has shown in [Vis96] that in the case of reflexive and transitive models, where well-foundedness is false and therefore can be expressed by a modal formula, the non-triviality of the fixpoint hierarchy is testified by the formula stating the existence of an infinite path alternately labelled with  $p$ ,  $\neg p$ ,  $p$ ,  $\neg p$ , etc.

In this paper we answer positively Lenzi's conjecture for the class of all transitive systems by giving an explicit syntactical translation of the full modal  $\mu$ -calculus into the alternation-free fragment. This result is first showed for finite transition systems and then generalized, by proving a finite model theorem, to all transitive systems. We also verify, again by giving an explicit syntactical translation, that if we add symmetry to transitivity all collapses to the purely modal fragment. Further, by adapting Arnold's proof for the general case, we show that the hierarchy remains strict over reflexive frames. By proving a finite model theorem for reflexive transition systems the corresponding result holds even on finite models.

In the next section we introduce the modal  $\mu$ -calculus and some additional, not standard, notions. In Section 3 we introduce evaluation games and show their relevance for the modal  $\mu$ -calculus. In Section 4 some finite model theorems are proved. In Sections 5 and 6 the collapse of the hierarchy over transitive-

---

<sup>4</sup> A complete proof of this fact, extended to the class of finite simple graphs (a class which contains - modulo bisimulation - the class of finite transitive graphs) can be found in [DL $\infty$ ].

symmetric and over transitive systems are proved. In Section 7 we prove the strictness of the hierarchy over reflexive transition systems.

### Related work

The question whether the modal  $\mu$ -calculus hierarchy collapses on special classes of transition systems has been addressed in various other works. A prominent subclass, coming from Gödel-Löb logic, is the class of transitive upward well-founded frames. As shown by Visser in [Vis05] and van Benthem in [vBe06] by using the de Jongh-Sambin fixpoint Theorem, the modal  $\mu$ -calculus collapses to its modal fragment. A direct proof of this result without using de Jongh-Sambin Theorem is given by the authors in [AF $\infty$ ]. In [tCFL07] ten Cate, Fontaine and Litak show that on the class of finite trees with the descendant relation the modal  $\mu$ -calculus collapses to the modal fragment. Concerning the hierarchy on transitive frames d'Agostino and Lenzi in [DL $\infty$ ] propose a different proof which uses Theorem 24 of this paper. Further, Dawar and Otto in [DO $\infty$ ] give a characterization of the bisimulation invariant fragment of Monadic Second Order Logic over transitive frames. From their result, by using the Finite Model Theorem for transitive frames (Theorem 15), the collapse of the modal  $\mu$ -calculus follows, too.

## 2 The propositional modal $\mu$ -calculus

In this section we introduce syntax, semantics of the modal  $\mu$ -calculus and the alternation depth hierarchy.

### 2.1 Syntax

The *language of the modal  $\mu$ -calculus*,  $\mathcal{L}_\mu$ , results by adding greatest and least fixpoint operators to propositional modal logic. More precisely, given a set  $\mathsf{P}$  of *propositional variables*, the collection  $\mathcal{L}_\mu$  of *modal  $\mu$ -formulae* (or simply  $\mu$ -formulae) is defined as follows:

$$\varphi ::= p \mid \sim p \mid \top \mid \perp \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid \diamond\varphi \mid \square\varphi \mid \mu x.\varphi \mid \nu x.\varphi$$

where  $p, x \in \mathsf{P}$  and  $x$  occurs only positively in  $\eta x.\varphi$  ( $\eta = \nu, \mu$ ), that is,  $\sim x$  is not a subformula of  $\varphi$ .  $\mathcal{L}_M$  denotes the pure modal fragment of  $\mathcal{L}_\mu$ .

The fixpoint operators  $\mu$  and  $\nu$  can be viewed as quantifiers. Therefore we use the standard terminology and notations as for quantifiers and, for instance,  $\mathsf{free}(\varphi)$  denotes the set of all propositional variables occurring free in  $\varphi$  and  $\mathsf{bound}(\varphi)$  those occurring bound. Further, we define  $\mathsf{var}(\varphi) = \mathsf{free}(\varphi) \cup \mathsf{bound}(\varphi)$ . If  $\psi$  is a subformula of  $\varphi$ , we write  $\psi \leq \varphi$ . We write  $\psi < \varphi$  when  $\psi$  is a proper subformula.  $\mathsf{sub}(\varphi)$  is the set of all subformulae of  $\varphi$ .

Let  $\varphi(x)$  and  $\psi$  be two  $\mu$ -formulae. The substitution of all occurrences of  $x$  with  $\psi$  in  $\varphi$  is denoted by  $\varphi[x/\psi]$  or sometimes simply  $\varphi(\psi)$ . Simultaneous

substitution of all  $x_i$  by  $\psi_i$  ( $i \in \{1, \dots, n\}$ ) is denoted by  $\varphi[x_1/\psi_1, \dots, x_n/\psi_n]$ . For serial substitution such as  $(\varphi[x_1/\psi_1])[x_2/\psi_2]$  we often omit the parentheses and write  $\varphi[x_1/\psi_1][x_2/\psi_2]$ .

*Remark 1.* Note, that if  $\varphi(x), \psi \in \mathcal{L}_\mu$  then  $\varphi[x/\psi]$  need not be a  $\mu$ -formula, for example, if we set  $\varphi \equiv \mu y.x$  and  $\psi \equiv \sim y$  then we have  $\varphi[x/\psi] \equiv \mu y. \sim y \notin \mathcal{L}_\mu$ . Nevertheless, in this paper, if nothing else mentioned, an expression like  $\varphi[x/\psi]$  will denote well defined  $\mu$ -formula. For a formal introduction of substitution we refer to Alberucci [Alb08].

The negation  $\neg\varphi$  of a  $\mu$ -formula  $\varphi$  is defined inductively such that  $\neg p \equiv \sim p$  and  $\neg(\sim p) \equiv p$ , by using de Morgan dualities for boolean connectives and the usual modal dualities for  $\diamond$  and  $\square$ . For  $\mu, \nu$  we define

$$\neg\mu x.\varphi(x) \equiv \nu x.\neg\varphi(x)[x/\neg x] \quad \text{and} \quad \neg\nu x.\varphi(x) \equiv \mu x.\neg\varphi(x)[x/\neg x].$$

As usual, we introduce implication  $\varphi \rightarrow \psi$  as  $\neg\varphi \vee \psi$  and equivalence  $\varphi \leftrightarrow \psi$  as  $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$ .

We say that a variable  $x \in \text{bound}(\varphi)$  is well-bounded in  $\varphi$  if no two distinct occurrences of fixpoint operators in  $\varphi$  bind  $x$ , and  $x$  occurs only once in  $\varphi$ . A propositional variable  $p$  is guarded in a formula  $\varphi \in \mathcal{L}_\mu$  if every occurrence of  $p$  in  $\varphi$  is in the scope of a modal operator. A formula  $\varphi$  of  $\mathcal{L}_\mu$  is said to be *guarded* if and only if for every subformula of  $\varphi$  of the form  $\eta x.\delta$ ,  $x$  is guarded in  $\delta$ . A formula  $\varphi$  of  $\mathcal{L}_\mu$  is said to be *well-named* if it is guarded and every  $x \in \text{bound}(\varphi)$  is well-bounded in  $\varphi$ . For all well-named  $\varphi$ , if  $x$  is bound in  $\varphi$  then there is exactly one subformula  $\eta x.\delta \leq \varphi$  which bounds  $x$ , this formula is denoted by  $\varphi_x$ . In the subsection of the semantics, by Lemma 2 we will see that any  $\mu$ -formula  $\varphi$  is equivalent to a well-named formula  $\text{wn}(\varphi)$ , therefore, if nothing else mentioned, we assume that all formulae are well-named.

If  $x \in \text{bound}(\varphi)$  and  $x$  is in the scope of a  $\diamond$  operator in  $\varphi_x$ , resp.  $\square$  operator, then we say that  $x$  is *weakly existential in  $\varphi$* , resp. *weakly universal in  $\varphi$* . If  $x \in \text{bound}(\varphi)$  and  $x$  is in the scope only of  $\diamond$  operators in  $\varphi_x$ , resp.  $\square$  operators, then we say that  $x$  is *existential in  $\varphi$* , resp. *universal in  $\varphi$* . Let  $\varphi(x)$  be a  $\mu$ -formula. If  $x$  is free and occurs only positively in  $\varphi$ , then we define  $\varphi^n(x)$  for all  $n$  inductively such that  $\varphi^1(x) = \varphi(x)$  and such that

$$\varphi^{k+1}(x) \equiv \varphi[x/\varphi^k(x)].$$

We define  $\varphi^n(\top) = \varphi^n[x/\top]$ , and analogously for  $\varphi^n(\perp)$ .

The *rank*,  $\text{rank}(\varphi)$ , of a formula  $\varphi$  is an ordinal number defined inductively as follows:

- $\text{rank}(p) = \text{rank}(\sim p) = 1$
- $\text{rank}(\Delta \alpha) = \text{rank}(\alpha) + 1$  where  $\Delta \in \{\square, \diamond\}$
- $\text{rank}(\alpha \circ \beta) = \max\{\text{rank}(\alpha), \text{rank}(\beta)\} + 1$  where  $\circ \in \{\wedge, \vee\}$
- $\text{rank}(\eta x.\alpha) = \sup\{\text{rank}(\alpha^n(x)) + 1 \ ; \ n \in \mathbb{N}\}$  where  $\eta \in \{\nu, \mu\}$ .

The fact that the definition of  $\text{rank}$  terminates is shown in the joint work with Krähenbühl [AK $\infty$ ] (see also [Alb08]). It is an easy exercise to show that for all formulae  $\varphi$  we have that  $\text{rank}(\varphi) = \text{rank}(\neg\varphi)$ .

The next lemma shows that wellnaming iterated formulae which are already well-named does not affect the rank. It follows by the fact that since  $\varphi$  is well-named the equivalent well-named formula for  $\varphi^n(\top)$  is given by simply renaming bound variables.

**Lemma 1** *For all well-named formulae  $\varphi$  such that  $x$  appears only positively and all  $n \in \mathbb{N}$  we have that*

$$\text{rank}(\varphi^n(\top)) = \text{rank}(\text{wn}(\varphi^n(\top))).$$

*Similarly for  $\perp$ .*

Given a  $\mu$ -formula  $\varphi$ , for all set of bound variables  $X \subseteq \text{bound}(\varphi)$ , the formula  $\varphi^{\text{free}(X)}$  is obtained from  $\varphi$  by eliminating all fixpoint operators binding a variable  $x \in X$  but leaving the previously bound variables  $x$  as a free occurrences. Further, if  $X = \{x_i, \dots, x_n\} \subseteq \text{bound}(\varphi)$  then we define

$$\varphi^{-X} \equiv \varphi^{\text{free}(X)}[x_1/\perp, \dots, x_n/\perp].$$

## 2.2 Semantics

The semantics of modal  $\mu$ -calculus is given by transition systems. A *transition system*  $\mathcal{T}$  is of the form  $(\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}})$  where  $\mathbf{S}$  is a set of *states*  $\rightarrow^{\mathcal{T}}$  is a binary relation on  $\mathbf{S}$  called the *accessibility relation* and  $\lambda : \mathbf{P} \rightarrow \wp(\mathbf{S})$  is a *valuation* for all propositional variables. A transition system  $\mathcal{T}$  with a distinguished state  $s$  is called a *pointed transition system* and denoted by  $(\mathcal{T}, s)$ .  $\mathbb{T}$  denotes the class of all pointed transition systems. Given any property  $P$ , by  $\mathbb{T}^P$  we denote the subclass of pointed transition systems satisfying the property  $P$ . In particular  $\mathbb{T}^r$  denotes all pointed reflexive transition systems,  $\mathbb{T}^{st}$  all pointed symmetric and transitive transition systems,  $\mathbb{T}^t$  all pointed transitive transition systems and  $\mathbb{T}^{rst}$  denotes all pointed transition systems where the accessibility relation is an equivalence relation, that is, it denotes the class of all **S5** models. Given any property  $P$ , with  $\mathbb{T}^{Pf}$  we denote the subclass of *finite* pointed transition systems satisfying the property  $P$ . For example,  $\mathbb{T}^{tf}$  denotes all *finite* pointed transition systems where the accessibility relation is transitive.

Let  $\lambda$  be a valuation,  $p$  a propositional variable and  $S'$  a subset of states  $\mathbf{S}$ ; we set for all propositional variables  $p'$

$$\lambda[p \mapsto S'](p') = \begin{cases} S' & \text{if } p' = p, \\ \lambda(p') & \text{otherwise.} \end{cases}$$

Given a transition system  $\mathcal{T} = (\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}})$ , then  $\mathcal{T}[p \mapsto S']$  denotes the transition system  $(\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}}[p \mapsto S'])$ . This notions are generalized straightforwardly to  $\lambda[x_1 \mapsto S_1, \dots, x_n \mapsto S_n]$  and  $\mathcal{T}[x_1 \mapsto S_1, \dots, x_n \mapsto S_n]$ . Given a transition

system  $\mathcal{T}$ , the *denotation of  $\varphi$  in  $\mathcal{T}$* ,  $\|\varphi\|_{\mathcal{T}}$ , that is, the set of states satisfying a formula  $\varphi$  is defined inductively on the structure of  $\varphi$ . Simultaneously for all transition systems we set

- $\|p\|_{\mathcal{T}} = \lambda(p)$  and  $\|\sim p\|_{\mathcal{T}} = \mathbf{S} - \lambda(p)$  for all  $p \in \mathbf{P}$ ,
- $\|\alpha \wedge \beta\|_{\mathcal{T}} = \|\alpha\|_{\mathcal{T}} \cap \|\beta\|_{\mathcal{T}}$ ,
- $\|\alpha \vee \beta\|_{\mathcal{T}} = \|\alpha\|_{\mathcal{T}} \cup \|\beta\|_{\mathcal{T}}$ ,
- $\|\Box\alpha\|_{\mathcal{T}} = \{s \in \mathbf{S} \mid \forall t((s \rightarrow^{\mathcal{T}} t) \Rightarrow t \in \|\alpha\|_{\mathcal{T}})\}$ ,
- $\|\Diamond\alpha\|_{\mathcal{T}} = \{s \in \mathbf{S} \mid \exists t((s \rightarrow^{\mathcal{T}} t) \wedge t \in \|\alpha\|_{\mathcal{T}})\}$ ,
- $\|\nu x.\alpha\|_{\mathcal{T}} = \bigcup\{S' \subseteq \mathbf{S} \mid S' \subseteq \|\alpha(x)\|_{\mathcal{T}[x \mapsto S']}\}$ , and
- $\|\mu x.\alpha\|_{\mathcal{T}} = \bigcap\{S' \subseteq \mathbf{S} \mid \|\alpha(x)\|_{\mathcal{T}[x \mapsto S']} \subseteq S'\}$ .

We say that a pointed transition system  $(\mathcal{T}, s)$  is a model of a  $\mu$ -formula if and only if  $s \in \|\varphi\|_{\mathcal{T}}$ . By  $\|\varphi\|$  we denote the class of all models of  $\varphi$  and by  $\|\varphi\|^P$  the class of all models of  $\varphi$  with property  $P$ . For a formula  $\varphi(x)$  and set of states  $S' \subseteq \mathbf{S}$  we sometimes write  $\|\varphi(S')\|_{\mathcal{T}}$  instead of  $\|\varphi(x)\|_{\mathcal{T}[x \mapsto S']}$ . When clear from the context we use  $\|\varphi(x)\|_{\mathcal{T}}$  for the function

$$\|\varphi(x)\|_{\mathcal{T}} : \begin{cases} \wp(\mathbf{S}) \rightarrow \wp(\mathbf{S}) \\ S' \mapsto \|\varphi(S')\|_{\mathcal{T}}. \end{cases}$$

By Tarski-Knaster Theorem, c.f. [Tar55],  $\|\nu x.\alpha(x)\|_{\mathcal{T}}$  is the greatest fixpoint and  $\|\mu x.\alpha(x)\|_{\mathcal{T}}$  the least fixpoint of the operator  $\|\alpha(x)\|_{\mathcal{T}}$ .

The next two lemmas state some basic properties of denotations. Their proofs are left to the reader.

**Lemma 2** *For all transition systems  $\mathcal{T} = (\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}})$  and all formulae  $\varphi$  we have that*

1.  $\|\neg\varphi\|_{\mathcal{T}} = \mathbf{S} - \|\varphi\|_{\mathcal{T}}$ ,
2.  $\|\eta x.\eta y.\varphi(x, y)\|_{\mathcal{T}} = \|\eta x.\varphi(x, x)\|_{\mathcal{T}}$ , where  $\eta \in \{\mu, \nu\}$ ,
3.  $\|\nu x.\varphi(x)\|_{\mathcal{T}} = \|\varphi(\top)\|_{\mathcal{T}}$ , if all occurrences of  $x$  are not guarded,
4.  $\|\mu x.\varphi(x)\|_{\mathcal{T}} = \|\varphi(\perp)\|_{\mathcal{T}}$ , if all occurrences of  $x$  are not guarded.
5. There is a well-named formula  $\text{wn}(\varphi)$  such that  $\|\varphi\|_{\mathcal{T}} = \|\text{wn}(\varphi)\|_{\mathcal{T}}$ .

**Lemma 3** *Given well-named formulae  $\varphi, \alpha, \alpha_i, \beta, \beta_i, \psi, \psi_i \in \mathcal{L}_{\mu}$ ,  $i \in \{1, \dots, k\}$ . For all transition systems  $\mathcal{T}$  the following holds:*

1. If  $\text{free}(\psi_i) \cap \text{bound}(\varphi) = \emptyset$  for all  $i \in \{1, \dots, k\}$  then

$$\|\varphi[x_1/\psi_1, \dots, x_k/\psi_k]\|_{\mathcal{T}} = \|\varphi\|_{\mathcal{T}[x_1 \mapsto \|\psi_1\|_{\mathcal{T}}, \dots, x_k \mapsto \|\psi_k\|_{\mathcal{T}}]}.$$

2. If  $\psi \leq \varphi$  and  $x_i \in \text{free}(\psi) \cap \text{bound}(\varphi)$ , with  $i = 1, \dots, k$ , then

$$\|\psi[x_1/\varphi_{x_1}, \dots, x_k/\varphi_{x_k}]\|_{\mathcal{T}} = \|\psi\|_{\mathcal{T}[x_1 \mapsto \|\varphi_{x_1}\|_{\mathcal{T}}, \dots, x_k \mapsto \|\varphi_{x_k}\|_{\mathcal{T}}]}.$$

3. If  $\text{free}(\psi_i) \cap \text{bound}(\alpha) = \text{free}(\psi_i) \cap \text{bound}(\beta) = \emptyset$  and  $x_i \in \text{free}(\alpha) \cap \text{free}(\beta)$  for all  $i \in \{1, \dots, k\}$  and for every transition system  $\mathcal{T}$  we have that

$$\|\alpha\|_{\mathcal{T}} = \|\beta\|_{\mathcal{T}}$$

then, for every transition system  $\mathcal{T}$  we have that

$$\|\alpha[x_1/\psi_1, \dots, x_k/\psi_k]\|_{\mathcal{T}} = \|\beta[x_1/\psi_1, \dots, x_k/\psi_k]\|_{\mathcal{T}}.$$

4. Let  $\text{free}(\alpha_i) \cap \text{bound}(\varphi) = \text{free}(\beta_i) \cap \text{bound}(\varphi) = \emptyset$  and let  $x_i \in \text{free}(\varphi)$  occur positively in  $\varphi$ , where  $i = 1, \dots, k$ . If for every transition system  $\mathcal{T}$  we have that

$$\|\alpha_i\|_{\mathcal{T}} \subseteq \|\beta_i\|_{\mathcal{T}}, \text{ for every } i \in \{1, \dots, k\},$$

then we have that

$$\|\varphi[x_1/\alpha_1, \dots, x_k/\alpha_k]\|_{\mathcal{T}} \subseteq \|\varphi[x_1/\beta_1, \dots, x_k/\beta_k]\|_{\mathcal{T}}.$$

Let  $\mathcal{T} = (\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda)$  be a transition system and  $s, s'$  two states in  $\mathbf{S}$ . A sequence  $s_0, s_1, \dots, s_n$  such that  $s_i \rightarrow^{\mathcal{T}} s_{i+1}$ ,  $s_0 = s$  and  $s_n = s'$  is a *path of length  $n$*  connecting  $s$  to  $s'$ . We say that  $s'$  is *reachable* from  $s$ . A subset  $S' \subseteq \mathbf{S}$  of the set of states is called a *strongly connected component* if for all  $s, s' \in S'$  we have that  $s'$  is reachable from  $s$ . For each  $s$  by  $\text{scc}(s)$  we denote the greatest strongly connected component which contains  $s$  if there is one and  $\text{scc}(s) = \emptyset$  if  $s$  is not contained in any strongly connected component. Note, that the notion  $\text{scc}(s)$  is well-defined. Given a pointed transition system  $(\mathcal{T}, s)$  and a state  $s'$  in it, we define the *depth of  $s'$* ,  $dp(s')$ , to be the length of the shortest path from  $s$  to  $s'$ . Since parts which are non connected to the point  $s$  will be irrelevant in the sequel we assume that all transition system are connected and, therefore, that  $dp(s')$  is defined for all  $s'$ .

### 2.3 The alternation depth hierarchy

Let  $\Phi \subseteq \mathcal{L}_{\mu}$ . For  $\eta \in \{\nu, \mu\}$ ,  $\eta(\Phi)$  is the smallest class of formulae such that:

- $\Phi, \neg\Phi \in \eta(\Phi)$ ;
- If  $\psi(x) \in \eta(\Phi)$  and  $x$  occurs only positively, then  $\eta x.\psi \in \eta(\Phi)$ ;
- If  $\psi, \varphi \in \eta(\Phi)$ , then  $\psi \wedge \varphi, \psi \vee \varphi, \diamond\psi, \square\psi \in \eta(\Phi)$ ;
- If  $\psi, \varphi \in \eta(\Phi)$  and  $\text{free}(\psi) \cap \text{bound}(\varphi) = \emptyset$  then  $\varphi[x/\psi] \in \eta(\Phi)$

With the help of this definition, we introduce the syntactical hierarchy for the modal  $\mu$ -calculus. For all  $n \in \mathbb{N}$ , we define the class of  $\mu$ -formulae  $\Sigma_n^{\mu}$  and  $\Pi_n^{\mu}$  inductively as follows:

- $\Sigma_0^{\mu} := \Pi_0^{\mu} := \mathcal{L}_{\mathbf{M}}$ ;
- $\Sigma_{n+1}^{\mu} = \mu(\Pi_n^{\mu})$ ;
- $\Pi_{n+1}^{\mu} = \nu(\Sigma_n^{\mu})$ .

$$\Delta_n^\mu := \Sigma_n^\mu \cap \Pi_n^\mu$$

The fixpoint alternation depth,  $\text{ad}$ , of a formula is the number of non-trivial nestings of alternating least and greatest fixpoints. Formally, the alternation depth of  $\varphi \in \mathcal{L}_\mu$  is given by

$$\text{ad}(\varphi) := \inf\{k : \varphi \in \Delta_{k+1}^\mu\}.$$

All  $\Sigma_n^\mu$  and  $\Pi_n^\mu$  form the *syntactical modal  $\mu$ -calculus hierarchy*, which is strict. The fixpoint alternation free fragment corresponds to the class  $\Delta_2^\mu$ .

**Lemma 4** *For all  $\mu$ -formulae  $\varphi$  there is a well-named formula  $\text{wn}(\varphi)$  such that for all  $\mathcal{T}$  we have  $\|\varphi\|_{\mathcal{T}} = \|\text{wn}(\varphi)\|_{\mathcal{T}}$  and  $\text{ad}(\varphi) = \text{ad}(\text{wn}(\varphi))$ .*

*Proof.* We have just to verify that the construction of  $\text{wn}(\varphi)$  given by parts 2 to 4 of Lemma 2 does not increase the alternation depth of the formula. But this is straightforward.  $\dashv$

Given Lemma 1 and Lemma 4, we can assume that  $\text{wn}$  is a function associating to every formula  $\varphi$  a well-named formula  $\text{wn}(\varphi)$  which has the same alternation depth and the same denotation in every transition system and such that, if  $\varphi$  is well-named, it also preserves the rank of  $\varphi^n(\top)$  and  $\varphi^n(\perp)$ , for every  $n$ .

The *semantical modal  $\mu$ -calculus hierarchy over  $\mathbb{T}$*  consists of all  $\Sigma_n^{\mu\mathbb{T}}$  and  $\Pi_n^{\mu\mathbb{T}}$ , which are classes of pointed transition systems defined inductively as follows:

$$\begin{aligned}\Sigma_n^{\mu\mathbb{T}} &= \{\|\varphi\| : \varphi \in \Sigma_n^\mu\} \\ \Pi_n^{\mu\mathbb{T}} &= \{\|\varphi\| : \varphi \in \Pi_n^\mu\}\end{aligned}$$

As usual, the ambiguous classes are defined by

$$\Delta_n^{\mu\mathbb{T}} := \Sigma_n^{\mu\mathbb{T}} \cap \Pi_n^{\mu\mathbb{T}}.$$

The *semantical modal  $\mu$ -calculus hierarchy over  $\mathbb{T}^P$* , for any property  $P$ , is defined analogously.

**Theorem 5** ([Bra98<sup>1</sup>, Bra98<sup>2</sup>]) *The semantical modal  $\mu$ -calculus hierarchy over  $\mathbb{T}$  is strict.*

From now on, when we write about the *modal  $\mu$ -calculus hierarchy*, we always mean the *semantical modal  $\mu$ -calculus hierarchy*.

*Example 1.* It is instructive to have a look at two typical  $\mu$ -formulae. The first formula express the property of “always eventually  $p$ ”

$$\nu x.(\mu y.(p \vee \diamond y)) \wedge \square x).$$

Indeed, it says that from any node of a model, we can reach a node where  $p$  holds. Since this formula is in  $\Pi_1^\mu$ , this kind of property can be expressed without any alternation. Moreover, it can be shown that cannot be reduced to a purely modal formula. The second formula defines the property of “there is a path where  $p$  holds infinitely often”

$$\nu x.\mu y.((p \vee \diamond y)) \wedge \diamond x).$$

It can be verified that the alternation is really needed, that is, that the class of models of this formula is in  $\Pi_2^{\mu\mathbb{T}} \setminus \Sigma_2^{\mu\mathbb{T}}$ .



### 3 Evaluation games for the modal $\mu$ -calculus

Evaluation games are a very important tool in the modal  $\mu$ -calculus and will play a crucial role in the sequel. After introducing some general notions for infinite games we define the evaluation games. The last subsection is about the concept of game formula.

#### 3.1 Parity games

Let  $V$  be a set. By  $V^*$  we denote the set of finite sequences on  $V$ , and by  $V^+$  we denote the set of nonempty sequences. Finally, by  $V^\omega$  we denote the set of infinite sequences over  $V$ .

A *game*  $\mathcal{G}$  is defined in terms of an arena  $A$  and a winning condition  $W$ . In our case an arena is simply a bi-partite graph  $A = \langle V_0, V_1, E \rangle$ , where  $V_0 \cap V_1 = \emptyset$  and the edge relation, or set of moves, is  $E \subseteq (V_0 \cup V_1) \times (V_0 \cup V_1)$ . Let  $V = V_0 \cup V_1$  be the set of vertices, or positions, of the arena. Given two vertices  $a, b \in V$ , we say that  $b$  is a successor of  $a$ , if  $(a, b) \in E$ . The set of all successors of  $a$  is sometimes denoted by  $aE$  or  $E(a)$ . We say that  $b$  is reachable from  $a$  if there are  $a_1, \dots, a_n \in V$  such that  $a_1 = a$ ,  $a_n = b$  and for every  $0 < i < n$ ,  $a_{i+1} \in a_i E$ .

A *play* in the arena  $A$  can be finite or infinite. In the former case, the play is a non empty finite path  $\pi = a_1 \dots a_n \in V^+$  such that for every  $0 < i < n$ ,  $a_{i+1} \in a_i E$  and  $a_n E = \emptyset$ . In the last case, the play consists in an infinite path  $\pi = a_1 \dots a_n \dots \in V^\omega$  with  $a_{i+1} \in a_i E$  for every  $i > 0$ . Thus a finite or infinite play in a game can be seen as the trace of a token moved on the arena by two Players, Player 0 and Player 1, in such a way that if the token is in position  $a \in V_i$ , then Player  $i$  has to choose a successor of  $a$  where to move the token.

The set of *winning conditions*  $W$  is a subset of  $V^\omega$ . Thus, given a game  $\mathcal{G} = (A, W)$  a play  $\pi$  is winning for Player 0 iff

1. if  $\pi$  is finite, then the last position  $a_n$  of the play is in  $V_1$ ,
2. if  $\pi$  is infinite, then it must be a member of  $W$ .

A play is winning for Player 1 if it is not winning for Player 0. In this framework we are interested in what is called a *parity winning condition*. That is, given a set of vertices  $V$ , we assume a coloring or ranking function  $\Omega : V \rightarrow \omega$  such that  $\Omega[V]$  is bounded. Then, the set  $W$  of winning conditions is defined as the set of all infinite sequences  $\pi$  such that the greatest priority appearing infinitely often in  $\Omega(\pi)$  is even.

Let  $A$  be an arena. A strategy for Player  $i$  is simply a function  $\sigma_i : V^*V_i \rightarrow V$ , with  $i = 1, 2$ . A prefix  $a_1 \dots a_n$  of a play is said to be *compatible* or *consistent* with  $\sigma_i$  iff for every  $j$  with  $1 \leq j < n$  and  $a_j \in V_i$ , it holds that  $\sigma_i(a_1 \dots a_j) = a_{j+1}$ . A finite or infinite play is compatible or consistent with  $\sigma_i$  if each of its prefix which is in  $V^*V_i$  is compatible with  $\sigma_i$ . The strategy  $\sigma_i$  is said to be a *winning strategy* for Player  $i$  on  $W$  if every play consistent with  $\sigma_i$  is winning for Player  $i$ . A position  $a \in V$  is winning for Player  $i$  in the parity game  $\mathcal{G}$  iff there is a strategy  $\sigma$  for Player  $i$  such that every play compatible with  $\sigma$  which

starts from  $a$  is winning for Player  $i$ . A winning strategy  $\sigma$  is called *memoryless* if  $\sigma(a_1 \dots a_n) = \sigma(b_1 \dots b_n)$ , when  $a_n = b_n$ . For parity games we have a memoryless determinacy result.

**Theorem 6 ([EJ91, Mos91])** *In a parity game, one of the Players has a memoryless winning strategy from each vertex.*

Having in mind this theorem, in the sequel we assume that all winning strategies are memoryless, that is, a winning strategy in a parity games for Player 0 is a function  $\sigma : V_0 \rightarrow V$ , analogously for Player 1.

### 3.2 Evaluation games for the modal $\mu$ -calculus

In this subsection we will see, given  $\varphi \in \mathcal{L}_\mu$  and a pointed transition system  $(\mathcal{T}, s_0)$  with  $\mathcal{T} = (\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}})$ , how to determine the corresponding parity game  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ , called also the *evaluation game of  $\varphi$  over  $(\mathcal{T}, s_0)$* .

The arena of  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  is the triple  $\langle V_0, V_1, E \rangle$  which is defined recursively such that

$$\langle \varphi, s_0 \rangle \in V$$

(remember that  $V = V_0 \cup V_1$ ) and such that if  $\langle \psi, s \rangle \in V$  then we distinguish the following cases:

- If  $\psi \equiv (\neg)p$  and  $p \in \text{free}(\varphi)$ . In this case we set  $E\langle \psi, s \rangle = \emptyset$  and

$$\langle \psi, s \rangle \in V_1 \text{ iff } \begin{cases} s \in \lambda^{\mathcal{T}}(\psi) & \text{if } \psi \equiv p \\ s \notin \lambda^{\mathcal{T}}(\psi) & \text{if } \psi \equiv \neg p. \end{cases}$$

- If  $\psi \equiv x$  and  $x \in \text{bound}(\varphi)$ . In this case we set

$$(\langle \psi, s \rangle, \langle \varphi_x, s \rangle) \in E$$

and we have

$$\langle \psi, s \rangle \in V_0 \text{ iff } x \text{ is a } \mu\text{-variable.}$$

- If  $\psi \equiv \alpha \wedge \beta$  then we have  $\langle \psi, s \rangle \in V_1$ , and if  $\psi \equiv \alpha \vee \beta$  then we have  $\langle \psi, s \rangle \in V_0$ . In both cases it holds that

$$(\langle \psi, s \rangle, \langle \alpha, s \rangle) \in E \text{ and } (\langle \psi, s \rangle, \langle \beta, s \rangle) \in E$$

- If  $\psi \equiv \Box \alpha$  then we have  $\langle \psi, s \rangle \in V_1$ , and if  $\psi \equiv \Diamond \alpha$  then we have  $\langle \psi, s \rangle \in V_0$ . In both cases it holds that

$$(\langle \psi, s \rangle, \langle \alpha, s' \rangle) \in E \text{ for all } s' \text{ such that } s \xrightarrow{\mathcal{T}} s'.$$

- If  $\psi \equiv \nu x. \alpha$  then we have  $\langle \psi, s \rangle \in V_1$ , and if  $\psi \equiv \mu x. \alpha$  then we have  $\langle \psi, s \rangle \in V_0$ . In both cases it holds that

$$(\langle \psi, s \rangle, \langle \alpha, s \rangle) \in E.$$

We complete the definition of the parity game  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  by defining the (partial) priority function  $\Omega : V \rightarrow \omega$ . The function is first defined on states of the form  $\langle \eta x. \delta, s \rangle \in V$ , where  $\eta \in \{\mu, \nu\}$ . In this case we have that:

$$\Omega(\langle \psi, s \rangle) = \begin{cases} \text{ad}(\eta x. \delta) & \text{if } \eta = \mu \text{ and } \text{ad}(\eta x. \delta) \text{ is odd, or} \\ & \eta = \nu \text{ and } \text{ad}(\eta x. \delta) \text{ is even;} \\ \text{ad}(\eta x. \delta) - 1 & \text{if } \eta = \mu \text{ and } \text{ad}(\eta x. \delta) \text{ is even, or} \\ & \eta = \nu \text{ and } \text{ad}(\eta x. \delta) \text{ is odd.} \end{cases}$$

For a state of the form  $\langle x, s \rangle$ , where  $x \in \text{bound}(\varphi)$ , we set

$$\Omega(\langle x, s \rangle) := \Omega(\langle \varphi_x, s \rangle).$$

For all the other states  $\langle \alpha, s \rangle$  we distinguish two cases. If there is a least formula  $\eta x. \delta \in \text{sub}(\varphi)$  such that  $\eta x. \delta > \alpha$  we set

$$\Omega(\langle \alpha, s \rangle) := \Omega(\langle \eta x. \delta, s \rangle).$$

If there is no such formula then we set

$$\Omega(\langle \alpha, s \rangle) = \begin{cases} \min\{\Omega(\eta x. \delta) : \eta x. \delta \leq \alpha\} & \text{if } \varphi \in \Delta_n^\mu, n > 1 \\ 1 & \text{if } \varphi \in \Delta_1^\mu. \end{cases}$$

It can easily be seen that if there is a formula  $\eta x. \delta > \alpha$  then there is also a least one. Therefore, the second case refers to subformulae  $\alpha$  of  $\varphi$  which can not be regenerated by a fixpoint application in a parity game. In the following we simply write  $\min \Omega$  and  $\max \Omega$  instead of  $\min\{\Omega(\langle \alpha, s \rangle) : \langle \alpha, s \rangle \in V\}$  and of  $\max\{\Omega(\langle \alpha, s \rangle) : \langle \alpha, s \rangle \in V\}$ .

Remember that if the play  $\pi$  is finite, Player 0 wins iff the last vertex of the play belongs to  $V_1$ , and if the play  $\pi$  is infinite, Player 0 wins iff the greatest priority appearing infinitely often even.

**Theorem 7 ([ES89])**  $(\mathcal{T}, s) \in \|\varphi\|$  iff Player 0 has a winning strategy for  $\mathcal{E}(\varphi, (\mathcal{T}, s))$ .

This result can be seen as the “game-theoretical version” of what is usually called the *Fundamental Theorem* of the semantic of the modal  $\mu$ -calculus. The proofs of the following Lemma is by unwinding the definitions of winning strategy.

**Lemma 8** Let  $\mathcal{T} = (\mathbf{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}})$  be a transition system and  $\varphi(x_1, \dots, x_k)$  be a formula where all  $x_i$  occurs positively. Let  $\sigma$  be a strategy for Player 0 in  $\mathcal{E}(\varphi(x_1, \dots, x_k), (\mathcal{T}, s))$ . Suppose that for all vertices of the form  $\langle x_i, s' \rangle$  which are reachable by  $\sigma$  we have that  $s' \in A_i \subseteq \mathbf{S}$ , with  $i = 1, \dots, k$ . Then  $\sigma$  can be converted into a winning strategy for Player 0 in  $\mathcal{E}(\varphi(x_1, \dots, x_k), (\mathcal{T}[x_1 \mapsto A_1, \dots, x_k \mapsto A_k], s))$ .

*Example 2.* Evaluation game  $\mathcal{E}(\nu x. \Box((p \vee \Box \perp) \wedge x), (\mathcal{T}, s_1))$ .  $\mathcal{T}$  is as in Figure 1, that is, it has states  $\{s_1, s_2, s_3\}$  and  $p$  holds in  $s_1$  and  $s_2$ , and the accessibility relation is as depicted in Figure 1.




Fig. 1.

In Figure 2, you find the arena of  $\mathcal{E}(\nu x.\Box((p \vee \Box\perp) \wedge x), (\mathcal{T}, s_1))$ . In order to simplify the picture we identified vertices of the form  $\langle \nu x.\Box((p \vee \Box\perp) \wedge x), s \rangle$  with the vertices of the form  $\langle \Box((p \vee \Box\perp) \wedge x), s \rangle$ . Note, that this does not change essentially the evaluation game. Further, the graph given by the non-dotted edges represents the part of the arena which can be reached by a play given the strategy of Player 0 where he chooses, if there is the possibility, the non-dotted instead of the dotted move. Note, that it is a winning strategy. It is left as an exercise to verify that  $\nu x.\Box((p \vee \Box\perp) \wedge x)$  is valid if for all reachable states in a transition system we have that either, the state is terminal, or,  $p$  holds in the state.




Fig. 2.

### 3.3 Game formulae

Given a parity game  $\mathcal{E}(\varphi, (\mathcal{T}, s))$  for a formula  $\varphi$  we define the *pointed game transition system*  $\mathcal{T}(\mathcal{E}(\varphi, (\mathcal{T}, s))) = ((\mathbb{S}, \rightarrow^{\mathcal{T}}, \lambda^{\mathcal{T}}), s_0)$  such that the states  $\mathbb{S}$  are the vertices  $V$  and the distinguished state  $s_0 = \langle \varphi, s \rangle$ , and such that the transition relation  $\rightarrow^{\mathcal{T}}$  is the edge relation  $E$  of the parity game. If  $\text{ad}(\varphi) = n$  then the valuation  $\lambda^{\mathcal{T}}$  is specified for the new propositional variables

$$\{c_i : 0 \leq i \leq n\} \cup \{d_i : 0 \leq i \leq n\}.$$

For all  $\psi \in \text{sub}(\varphi)$  we define our valuation for these propositional variables such that

$$\begin{aligned}\lambda^{\mathcal{T}}(d_i) &= \{\langle \psi, s \rangle : \langle \psi, s \rangle \in V_0 \text{ and } \Omega(\langle \psi, s \rangle) = i\} \quad \text{and} \\ \lambda^{\mathcal{T}}(c_i) &= \{\langle \psi, s \rangle : \langle \psi, s \rangle \in V_1 \text{ and } \Omega(\langle \psi, s \rangle) = i\}.\end{aligned}$$

In the following we introduce the *game formulae* and show that with them it is possible to test the existence of a winning strategy for Player 0 in an evaluation game.

**Definition 1.** For all  $n \geq 1$  we define the  $\Sigma_n^\mu$  game formula  $W_{\Sigma_n^\mu}$  such that:

$$W_{\Sigma_n^\mu} := \begin{cases} \nu x_{n-1} \cdot \nu x_{n-2} \cdot \dots \cdot \nu x_0 (\bigvee_{i=0}^{n-1} (d_i \wedge \Diamond x_i) \vee \bigvee_{i=0}^{n-1} (c_i \wedge \Box x_i)) & n \text{ even} \\ \mu x_n \cdot \nu x_{n-1} \cdot \dots \cdot \mu x_1 (\bigvee_{i=1}^n (d_i \wedge \Diamond x_i) \vee \bigvee_{i=1}^n (c_i \wedge \Box x_i)) & n \text{ odd} \end{cases}$$

The  $\Pi_n^\mu$  game formula  $W_{\Pi_n^\mu}$  is defined such that:

$$W_{\Pi_n^\mu} := \begin{cases} \nu x_n \cdot \mu x_{n-1} \cdot \dots \cdot \mu x_1 (\bigvee_{i=1}^n (d_i \wedge \Diamond x_i) \vee \bigvee_{i=1}^n (c_i \wedge \Box x_i)) & n \text{ even} \\ \nu x_{n-1} \cdot \mu x_{n-2} \cdot \dots \cdot \nu x_0 (\bigvee_{i=0}^{n-1} (d_i \wedge \Diamond x_i) \vee \bigvee_{i=0}^{n-1} (c_i \wedge \Box x_i)) & n \text{ odd} \end{cases}$$

For  $n = 0$  we define

$$W_{\Sigma_0^\mu} := W_{\Pi_0^\mu} := W_{\Sigma_1^\mu}.$$

It is clear from definition that for all  $n \geq 1$  we have that  $W_{\Sigma_n^\mu} \in \Sigma_n^\mu$  and  $W_{\Pi_n^\mu} \in \Pi_n^\mu$ .

**Proposition 9 ([EJ91, Wal02])** Let  $\mathcal{G}$  an arbitrary parity game such that  $\min \Omega \in \{0, 1\}$  and  $\max \Omega = n$ . We have that if  $n$  is even (resp. odd):

- (a) if  $\min \Omega = 0$  then Player 0 has a winning strategy for  $\mathcal{G}$  if and only if  $\mathcal{T}(\mathcal{G}) \in \|W_{\Pi_{n+1}^\mu}\|$  (resp.  $\mathcal{T}(\mathcal{G}) \in \|W_{\Sigma_{n+1}^\mu}\|$ ),
- (b) if  $\min \Omega = 1$  then Player 0 has a winning strategy for  $\mathcal{G}$  if and only if  $\mathcal{T}(\mathcal{G}) \in \|W_{\Pi_n^\mu}\|$  (resp.  $\mathcal{T}(\mathcal{G}) \in \|W_{\Sigma_n^\mu}\|$ )

From Proposition 9 and the definition of an evaluation game, it follows immediately that:

**Corollary 10** Let  $\varphi$  be a  $\Pi_n^\mu$ -formula (resp.  $\Sigma_n^\mu$ -formula) and let  $(\mathcal{T}, s)$  be an arbitrary pointed transition system. We have that Player 0 has a winning strategy for  $\mathcal{E}(\varphi, (\mathcal{T}, s))$  if and only if  $\mathcal{T}(\mathcal{E}(\varphi, (\mathcal{T}, s))) \in \|W_{\Pi_n^\mu}\|$  (resp.  $(\mathcal{T}, s) \in \|W_{\Sigma_n^\mu}\|$ ).

Therefore, by applying Proposition 7 and Corollary 10, we have the following result:

**Corollary 11** Let  $\varphi$  be a  $\Pi_n^\mu$ -formula (resp.  $\Sigma_n^\mu$ -formula) and let  $(\mathcal{T}, s)$  be an arbitrary pointed transition system. We have that

$$(\mathcal{T}, s) \in \|\varphi\| \text{ if and only if } \mathcal{T}(\mathcal{E}(\varphi, (\mathcal{T}, s))) \in \|W_{\Pi_n^\mu}\| \text{ (resp. } (\mathcal{T}, s) \in \|W_{\Sigma_n^\mu}\|).$$

## 4 Finite model theorems

In this section we prove finite model theorems for the modal  $\mu$ -calculus over the class of all reflexive and over the class of all transitive transition systems. Let us first state the well-known finite model theorem for general transition systems.

**Theorem 12 ([Koz88,ES89])** *For all modal  $\mu$ -formulae  $\varphi$  for which there is a transition system  $\mathcal{T}$  and a state  $s$  in  $\mathcal{T}$  such that  $s \in \|\varphi\|_{\mathcal{T}}$  there is a finite transition system  $\mathcal{T}^F$  and a state  $s^F$  such that  $s^F \in \|\varphi\|_{\mathcal{T}^F}$ .*

### 4.1 Finite model theorem for reflexive transition systems

Let  $\varphi$  be a  $\mu$ -formula. By induction on the structure of  $\varphi$  we define the formula  $\varphi_{ref}$  as follows:

- $(\sim)p_{ref} \equiv (\sim)p$ ,
- $(\alpha \circ \beta)_{ref} = \alpha_{ref} \circ \beta_{ref}$  where  $\circ \in \{\wedge, \vee\}$ ,
- $(\Box\alpha)_{ref} = \Box\alpha_{ref} \wedge \alpha_{ref}$ ,
- $(\Diamond\alpha)_{ref} = \Diamond\alpha_{ref} \vee \alpha_{ref}$ , and
- $(\eta x.\alpha)_{ref} = \eta x.\alpha_{ref}$  where  $\eta \in \{\mu, \nu\}$ .

The next Lemma is by induction on the structure of the formula.

**Lemma 13** *Let  $\mathcal{T}$  be a finite transition system and let  $\mathcal{T}^{ref}$  be its reflexive closure. For all  $\mu$ -formulae  $\varphi$  the following holds*

$$s \in \|\varphi_{ref}\|_{\mathcal{T}} \text{ if and only if } s \in \|\varphi\|_{\mathcal{T}^{ref}}.$$

With the help of this lemma we can easily prove the finite model property for reflexive transition systems.

**Theorem 14** *For all modal  $\mu$ -formulae  $\varphi$  for which there is a reflexive transition system  $\mathcal{T}$  and a state  $s$  in  $\mathcal{T}$  such that  $s \in \|\varphi\|_{\mathcal{T}}$  there is a finite reflexive transition system  $\mathcal{T}^F$  and a state  $s^F$  such that  $s^F \in \|\varphi\|_{\mathcal{T}^F}$ .*

*Proof.* Let  $\varphi$  be a  $\mu$ -formula and  $\mathcal{T}$  a reflexive transition system with a state  $s$  such that  $s \in \|\varphi\|_{\mathcal{T}}$ . Since  $\mathcal{T}$  is reflexive we have that  $\mathcal{T} = \mathcal{T}^{ref}$  and therefore by Lemma 13 we have that

$$s \in \|\varphi_{ref}\|_{\mathcal{T}}.$$

By the general Finite Model Theorem 12 we get that there is a finite transition system  $\mathcal{T}^F$  and a state  $s_F$  such that

$$s_F \in \|\varphi_{ref}\|_{\mathcal{T}^F}.$$

If we define  $\mathcal{T}^{Fref}$  to be the reflexive closure of  $\mathcal{T}^F$  by applying again Lemma 13 we get

$$s_F \in \|\varphi\|_{\mathcal{T}^{Fref}}$$

and we have found the finite reflexive model and a state in it satisfying  $\varphi$ .  $\dashv$

## 4.2 Finite model theorem for transitive transition systems

Let  $\varphi$  be a  $\mu$ -formula. By induction on the structure of  $\varphi$  we define the formula  $\varphi_{tr}$  as follows:

- $(\sim)p_{tr} \equiv (\sim)p$ ,
- $(\alpha \circ \beta)_{tr} = \alpha_{tr} \circ \beta_{tr}$  where  $\circ \in \{\wedge, \vee\}$ ,
- $(\Box\alpha)_{tr} = \nu x. \Box(\alpha_{tr} \wedge x)$ ,
- $(\Diamond\alpha)_{tr} = \mu x. \Diamond(\alpha_{tr} \vee x)$ , and
- $(\eta x. \alpha)_{tr} = \eta x. \alpha_{tr}$  where  $\eta \in \{\mu, \nu\}$ .

As in the reflexive case, the next Lemma is proved by induction on the structure of the formula.

**Lemma 15** *Let  $\mathcal{T}$  be a finite transition system and let  $\mathcal{T}^{tr}$  be its transitive closure. For all  $\mu$ -formulae  $\varphi$  the following holds*

$$s \in \|\varphi_{tr}\|_{\mathcal{T}} \quad \text{if and only if} \quad s \in \|\varphi\|_{\mathcal{T}^{tr}}.$$

By using Lemma 15, mutatis mutandis, the proof of the finite model property for transitive transition systems is exactly the same as for Theorem 14.

**Theorem 16** *For all modal  $\mu$ -formulae  $\varphi$  for which there is a transitive transition system  $\mathcal{T}$  and a state  $s$  in  $\mathcal{T}$  such that  $s \in \|\varphi\|_{\mathcal{T}}$  there is a finite transitive transition system  $\mathcal{T}^F$  and a state  $s^F$  such that  $s^F \in \|\varphi\|_{\mathcal{T}^F}$ .*

## 5 The hierarchy on transitive and symmetric transition systems

In this section, we prove the collapse of the semantical modal  $\mu$ -calculus hierarchy over  $\mathbb{T}^{st}$  to the purely modal fragment. Let us begin with the following easy lemma.

**Lemma 17** *Let  $\mathcal{T}$  be a transitive transition system and let  $s' \in \text{scc}(s)$ . For all  $\mu$ -formulae  $\varphi$  we have that*

$$s \in \|\Delta\varphi\|_{\mathcal{T}} \quad \text{if and only if} \quad s' \in \|\Delta\varphi\|_{\mathcal{T}}$$

where  $\Delta \in \{\Box, \Diamond\}$ .

*Proof.* Suppose that  $s \in \|\Box\varphi\|_{\mathcal{T}}$ . This is equivalent to the fact that for all  $s''$  such that  $s \rightarrow^{\mathcal{T}} s''$  we have that  $s'' \in \|\varphi\|_{\mathcal{T}}$ . On the other hand by definition of  $\text{scc}$  we have for all  $s' \in \text{scc}(s)$  that  $s \rightarrow^{\mathcal{T}} s'$  and  $s' \rightarrow^{\mathcal{T}} s$ . Therefore, for an arbitrary state  $s''$  by transitivity we have  $s \rightarrow^{\mathcal{T}} s''$  if and only if  $s' \rightarrow^{\mathcal{T}} s''$ . This implies that  $s \in \|\Box\varphi\|_{\mathcal{T}}$  if and only if for all  $s''$  such that  $s' \rightarrow^{\mathcal{T}} s''$  we have that  $s'' \in \|\varphi\|_{\mathcal{T}}$  which itself is equivalent to  $s' \in \|\Box\varphi\|_{\mathcal{T}}$ . The case for  $\Diamond$  is proved similarly.  $\dashv$

**Theorem 18** *Let  $\mathcal{T}$  be a transitive and symmetric transition system. We have that*

$$\|\nu x.\varphi(x)\|_{\mathcal{T}} = \|\varphi(\varphi(\top))\|_{\mathcal{T}}.$$

*Proof.* The  $\subseteq$  inclusion is clear. For the  $\supseteq$  inclusion, define  $A = \|\varphi(\varphi(\top))\|_{\mathcal{T}}$ ; by definition of greatest fixpoint it is enough to show that we have

$$A \subseteq \|\varphi(A)\|_{\mathcal{T}}. \quad (1)$$

First recall that we assume that  $\nu x.\varphi(x)$  is well-named. This means that in the formula  $\varphi(x)$  the variable  $x$  is in the scope of a modal operator and occurs only once in  $\varphi$ . Therefore, we can assume that  $\varphi$  is of the form  $\beta(\Delta \alpha(x))$  where  $\Delta \in \{\diamond, \square\}$ . Moreover  $x$  we have that  $\Delta \alpha(x)$  and  $\Delta \alpha(\varphi(x))$  occur only once in the formula tree of  $\varphi(\varphi(x))$ . Let  $s \in A$ , by Proposition 10 there is a winning strategy  $\sigma$  for Player 0 in the evaluation game  $\mathcal{E}(\varphi(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s))$ . Let  $\pi$  be an arbitrary play consistent with  $\sigma$ . If  $\pi$  reaches a vertex of the form  $\langle \Delta \alpha(x), s' \rangle$  then the same play reaches a vertex of the form  $\langle \Delta \alpha(\varphi(x)), s'' \rangle$ . Since  $\sigma$  is a winning strategy for Player 0 by Proposition 10 we have that

$$s'' \in \|\Delta \alpha(\varphi(x))\|_{\mathcal{T}[x \mapsto \mathbf{S}]} \quad \text{and} \quad s' \in \|\Delta \alpha(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}.$$

Since  $\mathcal{T}$  is transitive and symmetric it clearly holds that  $s'' \in \text{scc}(s')$  and, by applying Lemma 17, we have

$$s' \in \|\Delta \alpha(\varphi(x))\|_{\mathcal{T}[x \mapsto \mathbf{S}]}.$$

Hence, we have shown that for all plays  $\pi$  consistent with  $\sigma$ , if  $\pi$  reaches a vertex of the form  $\langle \Delta \alpha(x), s' \rangle$  then, by Theorem 7, there is a winning strategy for Player 0 in the evaluation game  $\mathcal{E}(\Delta \alpha(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s'))$ . A fortiori, this implies that if  $\pi$  reaches a vertex of the form  $\langle \Delta \alpha(x), s' \rangle$  then there is a winning strategy  $\sigma_{\langle \Delta \alpha(x), s' \rangle}$  for Player 0 in the evaluation game  $\mathcal{E}(\Delta \alpha(x), (\mathcal{T}[x \mapsto \|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}], s'))$ . Therefore, since  $\|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]} \subseteq \mathbf{S}$ , the strategy  $\sigma^*$  given by following  $\sigma$  but switching to the corresponding  $\sigma_{\langle \Delta \alpha(x), s' \rangle}$  when a position of the form  $\langle \Delta \alpha(x), s' \rangle$  is reached, is winning for Player 0 in  $\mathcal{E}(\varphi(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s))$ . Let  $B := \|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}$ . By construction of  $\sigma^*$  we have that for all vertices of the form  $\langle x, v \rangle$  which are reachable by  $\sigma^*$  it holds that  $v \in B$ . Then, by applying Lemma 8,  $\sigma^*$  can be converted into a winning strategy for Player 0 in  $\mathcal{E}(\varphi(\varphi(x)), (\mathcal{T}[x \mapsto B], s))$ . By Theorem 7, we have that

$$s \in \|\varphi(\varphi(B))\|_{\mathcal{T}}$$

which can be reformulated as  $s \in \|\varphi(\varphi(\varphi(\top)))\|_{\mathcal{T}}$  or  $s \in \|\varphi(A)\|_{\mathcal{T}}$ . Therefore, we have proved Equation 1 and completed the proof.  $\dashv$

**Definition 2.** *The syntactical translation  $(\cdot)^t : \mathcal{L}_{\mu} \rightarrow \mathcal{L}_M$  is defined recursively on the structure of the formula such that  $p^t = p$ ,  $\perp^t = \perp$  and  $\top^t = \top$ , such that it distributes over boolean and modal connectives, and such that*

$$(\mu x.\varphi)^t = (\text{wn}(\varphi(\varphi(\perp))))^t \quad \text{and} \quad (\nu x.\varphi)^t = (\text{wn}(\varphi(\varphi(\top))))^t.$$



Note that  $(\varphi)^t$  is defined via an application of  $(\cdot)^t$  either to a strict subformula  $\psi$  of  $\varphi$ , or to a formula whose rank, by Lemma 1, is strictly smaller than the rank of  $\varphi$ . Thus  $(\cdot)^t$  terminates and is well-defined.

The next corollary proves that on transitive and symmetric models, the semantical hierarchy of the  $\mu$ -calculus collapses to the class  $\Delta_1^{\mu\mathbb{T}^{st}}$ . Its proof goes by induction on the rank of a formula and uses Theorem 18.

**Corollary 19** *On transitive and symmetric transition systems we have that*

$$\|\varphi\|_{\mathcal{T}} = \|\varphi^t\|_{\mathcal{T}}.$$

*Example 3.* If we look at our example from Section 2, for “always eventually  $p$ ”, we have that

$$\|\nu x.(\mu y.(p \vee \diamond y)) \wedge \Box x\|^{\mathbb{T}^{st}} = \|(p \vee \diamond p) \wedge \Box(p \vee \diamond p)\|^{\mathbb{T}^{st}}$$

and for “there is a path where  $p$  holds infinitely often”, we have that

$$\begin{aligned} \|\nu x.\mu y.((p \vee \diamond y)) \wedge \diamond x\|^{\mathbb{T}^{st}} \\ = \\ \|(p \vee \diamond(p \wedge \diamond((p \vee \diamond(p \wedge \diamond \top)) \wedge \diamond \top))) \wedge \diamond((p \vee \diamond(p \wedge \diamond \top)) \wedge \diamond \top)\|^{\mathbb{T}^{st}}. \end{aligned}$$

*Remark 2.* Because the previous proof applies to any S5 model, that is, for every  $\mathcal{T} \in \mathbb{T}^{rst}$  we have that:

$$\|\varphi\|_{\mathcal{T}} = \|\varphi^t\|_{\mathcal{T}}$$

The fact that the modal  $\mu$ -calculus hierarchy for S5-models collapses to the pure modal fragment is indeed not surprising since for a S5-formula  $\varphi$  there are only finitely many formulae with the same propositional variables which are not equivalent over  $\mathbb{T}^{rst}$  and, therefore, it can easily be shown that for all  $\nu x.\varphi(x) \in \mathcal{L}_\mu$  there is a  $n \in \mathbb{N}$  such that  $\|\varphi^n(\top)\|^{\mathbb{T}^{rst}} = \|\nu x.\varphi\|^{\mathbb{T}^{rst}}$ . The existence of only finitely many non equivalent formulae follows from the fact that for all S5-formulae  $\varphi$  there is a conjunctive modal normal form  $\psi$  such that  $\psi \equiv \delta_1 \wedge \delta_2 \wedge \dots \wedge \delta_n$  where  $\delta \equiv \alpha \vee \Box\beta_1 \vee \Box\beta_2 \vee \dots \vee \Box\beta_n \vee \diamond\gamma_1 \vee \diamond\gamma_2 \vee \dots \vee \diamond\gamma_m$  and  $\alpha$ ,  $\beta_i$  and  $\gamma_j$  are propositional formulae<sup>5</sup>.

## 6 The hierarchy on transitive transition systems

We show that the modal  $\mu$ -calculus hierarchy over  $\mathbb{T}^t$  collapses to the alternation-free fragment. This is done in four parts starting from subsection two. First, any modal  $\mu$ -formula is reduced to a semantically equivalent formula  $\tau(\varphi)$  such that normalized strategies on evaluation games, which will be introduced in the third subsection, have certain nice properties. Then, we encode such normalized winning strategies in modal  $\mu$ -formulae and, finally, we show the collapse for finite transitive transition system and, by using the previously proved finite model theorem, generalize it to all transitive transition systems.

In the next subsection some technical notions like the one of unfolding a formula in a model are introduced and some properties are proved.

<sup>5</sup> Cf. Chapter 5 in [HC96].

### 6.1 Some technical preliminaries

Remember that we suppose all  $\mu$ -formulae well-named. First we introduce the unfolding of a formula which generalizes the one of closure of a formula, introduced by Kozen in [Koz83]. In order to do this, let  $\varphi$  and  $\psi$  be any  $\mu$ -formulae such that  $\{x_1, \dots, x_n\} = X \subseteq \text{bound}(\varphi)$ . The *unfolding of  $\psi$  in  $\varphi$  over  $X$* ,  $\text{unf}_\varphi^X(\psi)$ , is the formula defined recursively such that  $\text{unf}_\varphi^0(\psi) \equiv \psi$  and such that if  $X$  is of the form  $\{x_1, \dots, x_n\}$  then

$$\text{unf}_\varphi^X(\psi) \equiv \psi[x_1/\text{unf}_\varphi^{X^{-1}}(\varphi_{x_1}), \dots, x_n/\text{unf}_\varphi^{X^{-n}}(\varphi_{x_n})]$$

where  $X^{-i} = \{x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n\}$ . It can easy be seen that we have  $X \cap \text{free}(\text{unf}_\varphi^X(\psi)) = \emptyset$ .

In order to explain semantically the unfolding of a formula we introduce for each transition system  $\mathcal{T}$  the *transition system induced by  $\varphi$* ,  $\mathcal{T}^\varphi$ . For all  $x \in \text{bound}(\varphi)$  we define a natural number  $l(x)$  recursively such that  $l(x) = 0$  if  $\text{free}(\varphi_x) \cap \text{bound}(\varphi) = \emptyset$  and such that

$$l(x) = \max\{l(x_i) : x_i \in \text{free}(\varphi_x) \cap \text{bound}(\varphi)\} + 1$$

in the opposite case. For all transition systems  $\mathcal{T}$  with valuation  $\lambda$  we define new valuations  $\lambda^i$  and transition systems  $\mathcal{T}^i$  for all  $0 \leq i \leq \max\{l(x) : x \in \text{bound}(\varphi)\} =: N$  such that  $\lambda^0 = \lambda$  and  $\mathcal{T}^0 = \mathcal{T}$ , and such that  $\mathcal{T}^{k+1}$  is identical to  $\mathcal{T}^k$  except for the valuation  $\lambda^{k+1}$  which is defined as follows:

- $\lambda^{k+1}|_{\mathbb{P} \setminus \text{bound}(\varphi)} = \lambda^k|_{\mathbb{P} \setminus \text{bound}(\varphi)}$ ;
- if  $x \in \text{bound}(\varphi)$ :

$$\lambda^{k+1}(x) = \begin{cases} \lambda^k(x) & \text{if } l(x) \neq k + 1 \\ \|\varphi_x\|_{\mathcal{T}^k} & \text{if } l(x) = k + 1. \end{cases}$$

We define  $\mathcal{T}^\varphi$  to be  $\mathcal{T}^N$  and  $\lambda^\varphi = \lambda^N$ . Note, that if we have a formula  $\psi$  such that  $\text{free}(\psi) \cap \text{bound}(\varphi)$  is empty then, since the denotation of  $\varphi$  is independent of the valuation of the bound variables, we have  $\|\psi\|_{\mathcal{T}} = \|\psi\|_{\mathcal{T}^\varphi}$ . In particular, we have  $\|\varphi\|_{\mathcal{T}} = \|\varphi\|_{\mathcal{T}^\varphi}$ . Moreover note that for all  $x_i \in \text{bound}(\varphi)$  it holds that  $\lambda^\varphi(x_i) = \|\varphi_{x_i}\|_{\mathcal{T}^\varphi}$ .

**Lemma 20** *For all formulae  $\varphi$ , all subformulae  $\psi \leq \varphi$ , all  $X \subseteq \text{bound}(\varphi)$ , and all transition systems  $\mathcal{T}$  we have that*

$$\|\psi\|_{\mathcal{T}^\varphi} = \|\text{unf}_\varphi^X(\psi)\|_{\mathcal{T}^\varphi}.$$

*Proof.* By induction on the size of  $X$ . If  $X$  is empty, then by definition of unfolding we have that

$$\text{unf}_\varphi^X(\psi) \equiv \psi$$

and the claim is trivial. For the inductive step, suppose that  $X \cap \text{free}(\psi)$  is the set  $\{x_1, \dots, x_m\}$ . Hence, by definition we have

$$\text{unf}_\varphi^X(\psi) \equiv \psi[x_1/\text{unf}_\varphi^{X^{-1}}(\varphi_{x_1}), \dots, x_m/\text{unf}_\varphi^{X^{-m}}(\varphi_{x_m})].$$

Since  $\text{bound}(\psi) \cap \text{free}(\varphi_{x_i}) = \emptyset$  and  $\text{free}(\text{unf}_\varphi^{X^{-1}}(\varphi_{x_i})) \subseteq \text{free}(\varphi_{x_i})$  for all  $i$  we get that  $\text{bound}(\psi) \cap \text{free}(\text{unf}_\varphi^{X^{-1}}(\varphi_{x_i})) = \emptyset$ . Therefore, by induction hypothesis and Lemma 3.1 we get

$$\|\text{unf}_\varphi^X(\psi)\|_{\mathcal{T}^\varphi} = \|\psi\|_{\mathcal{T}^\varphi[x_1 \mapsto \|\varphi_{x_1}\|_{\mathcal{T}^\varphi}, \dots, x_m \mapsto \|\varphi_{x_m}\|_{\mathcal{T}^\varphi]}.$$

Since for all  $x_i$  we have that  $\lambda^\varphi(x_i) = \|\varphi_{x_i}\|_{\mathcal{T}^\varphi}$  we get

$$\|\text{unf}_\varphi^X(\psi)\|_{\mathcal{T}^\varphi} = \|\psi\|_{\mathcal{T}^\varphi}.$$

□

The previous lemma tells us that on the transition system induced by  $\varphi$  the denotation of any subformula of  $\varphi$  and the denotation of any of its unfolding over  $\varphi$  are the same.

Other usefull properties of  $\mathcal{T}^\varphi$  are summarized in the next lemma.

**Lemma 21** *Let  $\mathcal{T} = (\mathcal{S}, \rightarrow^\mathcal{T}, \lambda^\mathcal{T})$  be a transition system,  $\varphi$  any  $\mu$ -formula and  $\psi \leq \varphi$ . Then:*

1. For every  $X \subseteq \text{bound}(\varphi)$  we have

$$\|\psi^{\text{free}(X)}\|_{\mathcal{T}^\varphi} = \|\psi\|_{\mathcal{T}^\varphi}.$$

2. For every  $X_1, X_2 \subseteq \text{bound}(\varphi)$ , where  $X_1 \cap X_2 = \emptyset$ , we have

$$\|\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2} \psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi} = \|\psi\|_{\mathcal{T}^\varphi}.$$

3. For every  $X_1, X_2 \subseteq \text{bound}(\varphi)$ , where  $X_1 \cap X_2 = \emptyset$ , we have

$$\|\text{unf}_\varphi^{X_2} \psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi} = \|\psi\|_{\mathcal{T}^\varphi}.$$

*Proof.* Part 1. By Lemma 3.2 and since  $\|\varphi_x\|_{\mathcal{T}^\varphi} = \lambda^{\mathcal{T}^\varphi}(x)$  for all  $x \in \text{bound}(\varphi)$  we get

$$\|\psi^{\text{free}(X)}\|_{\mathcal{T}^\varphi} = \|\psi^{\text{free}(X)}[x_1/\varphi_{x_1}, \dots, x_n/\varphi_{x_n}]\|_{\mathcal{T}^\varphi}.$$

The proof ends with a straightforward induction on the structure of  $\psi$  proving that for all transition systems  $\mathcal{T}$  we have

$$\|\psi\|_{\mathcal{T}^\varphi} = \|\psi^{\text{free}(X)}[x_1/\varphi_{x_1}, \dots, x_n/\varphi_{x_n}]\|_{\mathcal{T}^\varphi}.$$

The only non trivial step is the one where  $\psi$  is of the form  $\eta x. \alpha$  ( $\eta \in \{\mu, \nu\}$ ). In this case, note that if any  $x_i$  appears free in  $\alpha$  then  $x$  appears only bound in  $\varphi_{x_i}$ .

Part 2. We prove the equation by induction on the size of  $X_2$ . If  $X_2$  is empty, the equation holds by the previous point. For the inductive step, given  $\{x_{i_1}, \dots, x_{i_k}\} = X_2 \cap \text{free}(\psi^{\text{free}(X_1)})$ , we have that by definition of  $\text{unf}$  the formula  $\|\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2} \psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi}$  is equal to

$$\|\psi^{\text{free}(X_1)}[x_{i_1}/\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_1}}(\varphi^{\text{free}(X_1)})_{x_{i_1}}, \dots, x_{i_k}/\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_k}}(\varphi^{\text{free}(X_1)})_{x_{i_k}}]\|_{\mathcal{T}^\varphi}.$$

Since  $\text{free}(\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_1}}(\varphi^{\text{free}(X_1)})_{x_{i_1}}) \subseteq \text{free}(\varphi^{\text{free}(X_1)})_{x_{i_1}}$  and since we have that  $\text{free}(\varphi^{\text{free}(X_1)})_{x_{i_1}} \cap \text{bound}(\psi^{\text{free}(X_1)}) = \emptyset$  we get

$$\text{free}(\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_1}}(\varphi^{\text{free}(X_1)})_{x_{i_1}}) \cap \text{bound}(\psi^{\text{free}(X_1)}) = \emptyset.$$

With Lemma 3.1 we get the equality with

$$\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi[x_{i_1} \mapsto \|\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_1}}(\varphi^{\text{free}(X_1)})_{x_{i_1}}\|_{\mathcal{T}^\varphi}, \dots, x_{i_k} \mapsto \|\text{unf}_{\varphi^{\text{free}(X_1)}}^{X_2^{-i_k}}(\varphi^{\text{free}(X_1)})_{x_{i_k}}\|_{\mathcal{T}^\varphi}]}$$

and by induction hypothesis this expression is equal to

$$\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi[x_{i_1} \mapsto \|\varphi_{x_{i_1}}\|_{\mathcal{T}^\varphi}, \dots, x_{i_k} \mapsto \|\varphi_{x_{i_k}}\|_{\mathcal{T}^\varphi}]}$$

Since in  $\mathcal{T}^\varphi$  we have that  $\lambda(x_{i_j}) = \|\varphi_{x_{i_j}}\|_{\mathcal{T}^\varphi}$  the last expression is equal to  $\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi}$ .

Part 3. Suppose  $\{x_{i_1}, \dots, x_{i_k}\} = X_2 \cap \text{free}(\psi^{\text{free}(X_1)})$ . Following the same argumentation as in part 2 we get that  $\|\text{unf}_{\varphi}^{X_2} \psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi}$  is equal to

$$\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi[x_{i_1} \mapsto \|\text{unf}_{\varphi}^{X_2^{-i_1}} \varphi_{x_{i_1}}\|_{\mathcal{T}^\varphi}, \dots, x_{i_k} \mapsto \|\text{unf}_{\varphi}^{X_2^{-i_k}} \varphi_{x_{i_k}}\|_{\mathcal{T}^\varphi}]}$$

With Lemma 20 we get the equality with

$$\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi[x_{i_1} \mapsto \|\varphi_{x_{i_1}}\|_{\mathcal{T}^\varphi}, \dots, x_{i_k} \mapsto \|\varphi_{x_{i_k}}\|_{\mathcal{T}^\varphi}]}$$

and because in  $\mathcal{T}^\varphi$  we have that  $\lambda(x_{i_j}) = \|\varphi_{x_{i_j}}\|_{\mathcal{T}^\varphi}$  the last expression is equal to  $\|\psi^{\text{free}(X_1)}\|_{\mathcal{T}^\varphi}$  which by part 1 is equal to  $\|\psi\|_{\mathcal{T}^\varphi}$ .  $\dashv$

**Lemma 22** *Let  $\varphi$  be a  $\mu$ -formula and  $\mathcal{T} = (\mathbb{S}, \rightarrow^T, \lambda^T)$  be a transition system. For all  $X \subseteq \text{bound}(\varphi)$ , all  $x_k \in \bar{X} = \text{bound}(\varphi) \setminus X$ , all  $\psi \leq \varphi$  and all  $x \notin \bar{X}$  we have that*

1.  $\|\text{unf}_{\varphi^{-\bar{X}}}^X \psi^{-\bar{X}}\|_{\mathcal{T}} \subseteq \|\text{unf}_{\varphi^{-\bar{X}^{-k}}}^{X \cup \{x_k\}} \psi^{-\bar{X}^{-k}}\|_{\mathcal{T}}$ ,
2.  $\|\text{unf}_{\varphi^{-\bar{X}}}^X \psi^{-\bar{X}}\|_{\mathcal{T}} \subseteq \|\text{unf}_{\varphi}^{\text{bound}(\varphi)} \psi\|_{\mathcal{T}}$ ,
3.  $\|\text{unf}_{\varphi^{-\bar{X}}}^X (\varphi^{-\bar{X}})_x\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}$ ,
4.  $\|\psi^{-\bar{X}}\|_{\mathcal{T}^{\varphi-X}} \subseteq \|\psi\|_{\mathcal{T}^\varphi}$ .

*Proof.* Suppose  $\sigma$  is a winning strategy for Player 0 in  $\mathcal{E}(\text{unf}_{\varphi^{-\bar{X}}}^X \psi^{-\bar{X}}, (\mathcal{T}, s))$ . By definition, any winning play for Player 0 starting from  $\langle \text{unf}_{\varphi^{-\bar{X}}}^X \psi^{-\bar{X}}, s \rangle$  and compatible with  $\sigma$  do not reach a position of type  $\langle \perp, s' \rangle$ . Thus, this strategy determines a winning strategy for Player 0 in  $\mathcal{E}(\text{unf}_{\varphi^{-\bar{X}^{-k}}}^{X \cup \{x_k\}} \psi^{-\bar{X}^{-k}}, (\mathcal{T}, s))$ . Part 1 is then obtained by applying Theorem 7. Part 2 follows by a finite reiteration of part 1. In order to obtain part 3 just apply Lemma 20 to part 2 and note that, since  $x \notin \bar{X}$ ,  $(\varphi^{-\bar{X}})_x \equiv (\varphi_x)^{-\bar{X}}$ . Part 4 is also a consequence of an application of Lemma 20 to part 2.  $\dashv$

## 6.2 A first reduction

We begin with a Lemma whose proof is standard.

**Lemma 23** *Let  $\mathcal{T}$  be a transitive transition system and let  $s, s'$  be two states such that  $s \rightarrow^{\mathcal{T}} s'$ . For all  $\mu$ -formulae  $\varphi$  we have that*

$$\begin{aligned} s \in \|\Box\varphi\|_{\mathcal{T}} &\implies s' \in \|\Box\varphi\|_{\mathcal{T}} \quad \text{and} \\ s' \in \|\Diamond\varphi\|_{\mathcal{T}} &\implies s \in \|\Diamond\varphi\|_{\mathcal{T}}. \end{aligned}$$

**Theorem 24** *Let  $\mathcal{T}$  be a transitive transition system and let  $\nu x.\varphi(x)$  be a formula such that  $x$  is weakly universal. We have that*

$$\|\nu x.\varphi(x)\|_{\mathcal{T}} = \|\varphi(\top)\|_{\mathcal{T}}.$$

*Proof.* The  $\subseteq$  inclusion is clear. For the  $\supseteq$  inclusion, define  $A = \|\varphi(\top)\|_{\mathcal{T}}$ ; by definition of greatest fixpoint it is enough to show that we have

$$A \subseteq \|\varphi(A)\|_{\mathcal{T}}. \quad (2)$$

First, recall that we assume that  $\nu x.\varphi(x)$  is well-named. This means that in the formula  $\varphi(x)$  the variable  $x$  is in the scope of a modal operator and, therefore, we can assume that  $\varphi$  is of the form  $\beta(\Box\alpha(x))$ . Moreover  $x$  occurs only once in  $\varphi$ . This implies that  $\Box\alpha(x)$  and  $\Box\alpha(\varphi(x))$  occur only once in the formula tree of  $\varphi(\varphi(x))$ . Let  $s \in A$ , by Theorem 7 there is a winning strategy  $\sigma$  for Player 0 in the evaluation game  $\mathcal{E}(\varphi(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s))$ . Let  $\pi$  be an arbitrary play consistent with  $\sigma$ . If  $\pi$  reaches a vertex of the form  $\langle \Box\alpha(x), s' \rangle$  then the same play reaches a vertex of the form  $\langle \Box\alpha(\varphi(x)), s'' \rangle$ , with  $\Box\alpha(x) \leq \Box\alpha(\varphi(x))$  and  $s'$  reachable from  $s''$  in  $\mathcal{T}[x \mapsto \mathbf{S}]$ . Since  $\sigma$  is a winning strategy for Player 0 by Proposition 10 we have that

$$s'' \in \|\Box\alpha(\varphi(x))\|_{\mathcal{T}[x \mapsto \mathbf{S}]} \quad \text{and} \quad s' \in \|\Box\alpha(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}.$$

Since  $\mathcal{T}[x \mapsto \mathbf{S}]$  is transitive we have that  $s'' \rightarrow^{\mathcal{T}[x \mapsto \mathbf{S}]} s'$  and, by applying Lemma 23, we have

$$s' \in \|\Box\alpha(\varphi(x))\|_{\mathcal{T}[x \mapsto \mathbf{S}]}.$$

Hence, we have shown that for all plays  $\pi$  consistent with  $\sigma$ , if  $\pi$  reaches a vertex of the form  $\langle \Box\alpha(x), s' \rangle$  then, by Theorem 7, there is a winning strategy for Player 0 in the evaluation game  $\mathcal{E}(\Box\alpha(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s'))$ . A fortiori, this implies that if  $\pi$  reaches a vertex of the form  $\langle \Box\alpha(x), s' \rangle$  then there is a winning strategy  $\sigma_{\langle \Box\alpha(x), s' \rangle}$  for Player 0 in the evaluation game  $\mathcal{E}(\Box\alpha(x), (\mathcal{T}[x \mapsto \|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}], s'))$ . Therefore, since  $\|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]} \subseteq \mathbf{S}$ , the strategy  $\sigma^*$  given by following  $\sigma$  but switching to the corresponding  $\sigma_{\langle \Box\alpha(x), s' \rangle}$  when a position of the form  $\langle \Box\alpha(x), s' \rangle$  is reached, is winning for Player 0 in  $\mathcal{E}(\varphi(\varphi(x)), (\mathcal{T}[x \mapsto \mathbf{S}], s))$ . Let  $B := \|\varphi(x)\|_{\mathcal{T}[x \mapsto \mathbf{S}]}$ . By construction of  $\sigma^*$  we have that for all vertices of the form  $\langle x, v \rangle$  which are reachable by  $\sigma^*$  it holds that  $v \in B$ . Then, by

applying Lemma 8,  $\sigma^*$  can be converted into a winning strategy for Player 0 in  $\mathcal{E}(\varphi(\varphi(x), (\mathcal{T}[x \mapsto B], s)))$ . By Theorem 7, we have that

$$s \in \|\varphi(\varphi(B))\|_{\mathcal{T}[x \mapsto s]}$$

which can be reformulated as  $s \in \|\varphi(\varphi(\varphi(\top)))\|_{\mathcal{T}}$  or  $s \in \|\varphi(A)\|_{\mathcal{T}}$ . Therefore, we have proved Equation 2 and completed the proof.  $\dashv$

**Definition 3.** *The syntactical translation  $\tau : \mathcal{L}_\mu \rightarrow \mathcal{L}_\mu$  is defined recursively on the structure of the formula such that  $\tau(p) = p$ ,  $\tau(\neg p) = \neg p$ ,  $\tau(\perp) = \perp$  and  $\tau(\top) = \top$ , such that it distributes over boolean and modal connectives, and such that*

- $\tau(\mu x.\varphi) = \tau(\text{wn}(\varphi(\varphi(\perp))))$ ,  $x$  is weakly existential in  $\varphi$
- $\tau(\mu x.\varphi) = \text{wn}(\mu x.\tau(\varphi))$ ,  $x$  is universal in  $\varphi$
- $\tau(\nu x.\varphi) = \tau(\text{wn}(\varphi(\varphi(\top))))$ ,  $x$  is weakly universal in  $\varphi$
- $\tau(\nu x.\varphi) = \text{wn}(\nu x.\tau(\varphi))$ ,  $x$  is existential in  $\varphi$ .

First, note that in each defining clause  $\tau(\varphi)$  is defined via an application of  $\tau$  to a formula whose rank, by Lemma 1, is strictly smaller than the rank of  $\varphi$ . Thus  $\tau$  terminates and is well-defined. Note also, that it can be proved by induction on the structure of  $\varphi$  that all variables which are existential (resp. universal) in  $\varphi$  are weakly existential (resp. universal) in  $\tau(\varphi)$  and that therefore for all  $\mu x.\alpha \leq \tau(\varphi)$  we have that  $x$  is weakly universal and for all  $\nu x.\alpha \leq \tau(\varphi)$  we have that  $x$  is weakly existential.

**Corollary 25** *On transitive transition systems we have that*

$$\|\varphi\|_{\mathcal{T}} = \|\tau(\varphi)\|_{\mathcal{T}}.$$

*Proof.* By induction on  $\text{rank}(\varphi)$ . If  $\text{rank}(\varphi) = 1$  or  $\text{rank}(\varphi)$  is a successor ordinal the proof is straightforward. If  $\text{rank}(\varphi)$  is a limit ordinal then  $\varphi$  is of the form  $\eta x.\alpha$ . We distinguish four cases. If  $\varphi$  is of the form  $\nu x.\alpha$  and  $x$  is existential in  $\varphi$  the induction step is straightforward. Similarly for  $\varphi$  of the form  $\mu x.\alpha$  and  $x$  is universal in  $\varphi$ . If  $\varphi$  is of the form  $\nu x.\alpha$  and  $x$  is in the scope of a  $\square$  in  $\varphi$  the induction step follows from Theorem 24 and Lemma 4. In the third case, if  $\varphi$  is of the form  $\mu x.\alpha$  and  $x$  is in the scope of a  $\diamond$  in  $\varphi$  then  $\neg\varphi$  is of the form  $\nu x.\neg\alpha[x/\neg x]$  and  $x$  is in the scope of a  $\square$  in  $\neg\varphi$ . Since in this case  $\text{rank}(\varphi) = \text{rank}(\neg\varphi)$  we can apply the induction step as in the third case.  $\dashv$

### 6.3 Normalizing the winning strategies

Let  $\mathcal{T}$  be a transitive transition system and  $\varphi$  a  $\mu$ -formula. Consider an arbitrary (memoryless) strategy  $\sigma$  for Player 0, not necessarily winning. We define the restriction of  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  on  $\sigma$ , denoted by  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$ , as follows:

- The set of positions  $V|_\sigma$  of the restriction is given by all nodes which are the positions of some play compatible with  $\sigma$  starting from position  $\langle \varphi, s_0 \rangle$ ,

- The arena of  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  is the triple  $\langle V_0|_\sigma, V_1|_\sigma, E|_\sigma \rangle$  where:
  1.  $V_0|_\sigma = \emptyset$ ,
  2.  $V_1|_\sigma = V|_\sigma$ ,
  3. if  $\langle \psi, s \rangle \in V|_\sigma \cap V_1$  then  $E|_\sigma(\langle \psi, s \rangle) = E(\langle \psi, s \rangle)$ , and
  4. if  $\langle \psi, s \rangle \in V|_\sigma \cap V_0$  then  $E|_\sigma(\langle \psi, s \rangle) = \{\sigma(\langle \psi, s \rangle)\}$ .
- The ranking function  $\Omega|_\sigma$  is given by the restriction of  $\Omega$  on  $V|_\sigma$ .

Note, that if  $\mathcal{T}$  is finite then  $V|_\sigma$  is finite, too. We have that in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  the only Player who can move is Player 1. This can be done because the moves for Player 0 are already completely determined by the (memoryless) strategy  $\sigma$ . Clearly, any play in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  is a play in  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  compatible with  $\sigma$ . We say that a play  $\pi$  in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  is winning for Player 0 if and only if the play  $\pi$  is winning for Player 0 in  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . If  $\sigma$  is a winning strategy for Player 0 then any play in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  is winning for Player 0.

*Example 4.* Look at the arena depicted in Example 2. The non-dotted part of the picture represents the arena of a restricted evaluation game.

**Definition 4.** Let  $\mathcal{T}$  be a finite transitive transition system and  $\varphi$  a  $\mu$ -formula. Suppose there is a winning strategy  $\sigma$  for Player 0 in the parity game  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . Then, for every position  $\langle \psi, s \rangle$  of  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$ , we define a measure  $d(\langle \psi, s \rangle)$ . We distinguish two cases in the definition, depending on whether the strongly connected component  $\text{scc}(\langle \psi, s \rangle)$  of  $\langle \psi, s \rangle$  in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  is empty or not:

1.  $\text{scc}(\langle \psi, s \rangle) = \emptyset$  :

$$d(\langle \psi, s \rangle) = \begin{cases} 0 & \text{if } E|_\sigma(\langle \psi, s \rangle) = \emptyset \\ \max\{d(\langle \phi, s' \rangle) : \langle \phi, s' \rangle \in E|_\sigma(\langle \psi, s \rangle)\} + 1 & \text{else} \end{cases}$$

2.  $\text{scc}(\psi, s) \neq \emptyset$  :

$$d(\langle \psi, s \rangle) = 0 \quad \text{if} \quad \bigcup \{E|_\sigma(\langle \alpha, s \rangle) : \langle \alpha, s \rangle \in \text{scc}(\psi, s)\} \setminus \text{scc}(\psi, s) = \emptyset,$$

else

$$d(\langle \psi, s \rangle) = \max\{d(\langle \phi, s' \rangle) : \langle \phi, s' \rangle \notin \text{scc}(\langle \psi, s \rangle) \text{ and exists } \langle \xi, s'' \rangle \in \text{scc}(\langle \psi, s \rangle) \text{ with } \langle \phi, s' \rangle \in E|_\sigma(\langle \xi, s'' \rangle)\} + 1.$$

For all finite transition systems  $d$  is a well-defined measure. Indeed, if we have a finite transition system we obviously have a finite arena which can be collapsed to a finite and well-founded graph by identifying all vertices in the arena which are in the same strongly connected component. It is clear that on finite and well-founded graphs  $d$  is well-defined. By noting that on the original arena the measure of a vertex corresponds to its measure of the collapsed arena we get that  $d$  is well-defined.

**Lemma 26** Let  $\mathcal{T}$  be a finite transitive transition system and  $\varphi \in \Sigma_2^\mu$ . Suppose there is a winning strategy  $\sigma$  for Player 0 in the parity game  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . If  $y \in \text{bound}(\varphi)$  is a  $\mu$ -variable, then for every position  $\langle y, s \rangle \in V|_\sigma$ , we have that  $\text{scc}(\langle y, s \rangle) = \emptyset$ .

*Proof.* If  $\text{scc}(\langle y, s \rangle) \neq \emptyset$  then Player 1 can determine a play  $\pi$  in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  where  $\langle y, s \rangle$  occurs infinitely often, since in  $\mathcal{E}|_\sigma(\varphi, (\mathcal{T}, s_0))$  only Player 1 moves and therefore can stay as long as he wants in a strongly connected component. Remember that  $\varphi \in \Sigma_2^\mu$ . Thus, there is no  $\nu$ -variable free in  $\varphi_y$ . Moreover, if  $y \in \text{free}(\varphi_x)$ , where  $x$  is an arbitrary  $\nu$ -variable, we have that  $\Omega(\langle y, s \rangle)$  is strictly greater than the priorities of  $x$  and  $\varphi_x$  positions. Therefore,  $\pi$  is winning for Player 1. But since  $\pi$  is compatible with  $\sigma$ , the play must be winning for Player 0, too. A contradiction.  $\dashv$

Given the restriction of  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  on a winning strategy  $\sigma$  and the measure  $d$  on  $V|_\sigma$  we define the *normalization* of  $\sigma$ , denoted by  $\sigma^N$ , as follows:

- For all positions  $\langle \diamond\beta, s' \rangle \in V|_\sigma$  we have that

$$\sigma^N(\langle \diamond\beta, s' \rangle) = \sigma(\langle \diamond\beta, s' \rangle),$$

if

$$d(\sigma(\langle \diamond\beta, s' \rangle)) = \min\{d(\langle \beta, \bar{s} \rangle) : \langle \beta, \bar{s} \rangle \in E(\langle \diamond\beta, s' \rangle) \text{ reachable from } \langle \diamond\beta, s' \rangle \text{ in } V|_\sigma\}$$

else

$$\sigma^N(\langle \diamond\beta, s' \rangle) = \langle \beta, s'' \rangle,$$

where  $\langle \beta, s'' \rangle \in E(\langle \diamond\beta, s' \rangle)$  is a vertex reachable from  $\langle \diamond\beta, s' \rangle$  in  $V|_\sigma$  such that

$$d(\langle \beta, s'' \rangle) = \min\{d(\langle \beta, \bar{s} \rangle) : \langle \beta, \bar{s} \rangle \in E(\langle \diamond\beta, s' \rangle) \text{ reachable from } \langle \diamond\beta, s' \rangle \text{ in } V|_\sigma\}.$$

- If  $\psi$  is not of the form  $\diamond\beta$  then we simply set  $\sigma^N(\langle \psi, s \rangle) = \sigma(\langle \psi, s \rangle)$ .

Intuitively, given a winning strategy  $\sigma$  for Player 0 on  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ , the normalized strategy  $\sigma^N$  for Player 0 is given by adapting  $\sigma$  such that for all vertexes of the form  $\langle \diamond\beta, s' \rangle$  Player 0 moves to a vertex  $\langle \beta, s'' \rangle$  whose measure is the minimal measure of all positions of the type  $\langle \beta, \bar{s} \rangle$  reachable from  $\langle \diamond\beta, s' \rangle$  which are still winning in  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . We have the following lemma.

**Lemma 27** *Let  $\mathcal{T}$  be a finite transitive transition system. If  $\sigma$  is a winning strategy for Player 0 on  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  then  $\sigma^N$  is a winning strategy for Player 0 on  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ , too.*

*Proof.* First we prove the following claim:

**Claim :**  $E|_\sigma$  and  $E|_{\sigma^N}$  coincide on every non empty scc of  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ .

The proof of the claim goes as follow. If there is no position of the form  $\langle \diamond\beta, s \rangle$  in a scc of  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ , the claim is trivially verified. Consider now an arbitrary scc( $\langle \diamond\beta, s \rangle$ ) of  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ . Let  $\langle \psi, t \rangle \in \text{scc}(\langle \diamond\beta, s \rangle)$ , in order to prove the claim we have to show that  $E|_{\sigma^N}(\langle \psi, t \rangle) = E|_\sigma(\langle \psi, t \rangle)$ .

- (a) If  $\psi$  is not of the form  $\diamond\alpha$ , then  $E|_{\sigma^N}(\langle \psi, t \rangle) = E|_\sigma(\langle \psi, t \rangle)$ .



- (b) For the case where  $\psi = \diamond\alpha$  then suppose that  $E|_{\sigma^N}(\langle\psi, t\rangle) \neq E|_{\sigma}(\langle\psi, t\rangle)$  and that  $E|_{\sigma^N}(\langle\psi, t\rangle) = \{\langle\alpha, t'\rangle\}$ . Note, that by construction of  $\sigma^N$  the position  $\langle\alpha, t'\rangle$  is the only successor of  $\langle\psi, t\rangle$ . Since  $E|_{\sigma^N}(\langle\psi, t\rangle) \neq E|_{\sigma}(\langle\psi, t\rangle)$  it must hold that

$$d(\langle\alpha, t'\rangle) < d(\langle\psi, t\rangle) \quad (3)$$

where  $d$  is the depth defined on  $E|_{\sigma}(\varphi, (\mathcal{T}, s_0))$ . Since  $\text{scc}(\langle\diamond\alpha, t\rangle) \neq \emptyset$  and since  $\langle\alpha, t'\rangle$  is the only position reachable in one step from  $\langle\diamond\alpha, t\rangle$  we have that  $\langle\alpha, t'\rangle \in \text{scc}(\langle\diamond\alpha, t\rangle)$  and therefore that  $\langle\diamond\alpha, t\rangle$  is reachable from  $\langle\alpha, t'\rangle$  in  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ . Since reachability in  $E|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$  implies reachability in  $E|_{\sigma}(\varphi, (\mathcal{T}, s_0))$  we can infer that  $d(\langle\alpha, t'\rangle) \geq d(\langle\diamond\alpha, t\rangle)$ , where  $d$  is the depth defined on  $E|_{\sigma}(\varphi, (\mathcal{T}, s_0))$ . This is a contradiction to point 3 and therefore the claim is proved.

Consider an arbitrary play  $\pi$  in the graph of  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ . If  $\pi$  is finite, then by construction of the normalized arena the play is winning for Player 0. If  $\pi$  is infinite then from a certain position, say  $\langle\alpha, t\rangle$ , we are in a  $\text{scc}$  of  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$ . But then by the previous claim after  $\langle\alpha, t\rangle$  the strategies of  $\sigma$  and  $\sigma^N$  coincide. Since by construction of  $\sigma^N$  the position  $\langle\alpha, t\rangle$  is winning in  $\mathcal{E}|_{\sigma}(\varphi, (\mathcal{T}, s_0))$  the highest priority appearing infinitely often in  $\pi$  often must be even and, therefore  $\pi$  is a winning play in  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$  for Player 0.  $\dashv$

In the next lemma we prove that, when considering  $\Sigma_2^\mu$ -formulae, normalized strategies have a nice and very usefull property.

**Lemma 28** *Let  $\mathcal{T}$  be a finite transitive transition system and  $\varphi \in \Sigma_2^\mu$  such that all  $\nu$ -variables are weakly existential. Let  $\sigma^N$  be a normalized winning strategy for Player 0 on  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . Consider a position  $\langle x, s_1 \rangle$  in  $\mathcal{E}|_{\sigma^N}(\varphi, (\mathcal{T}, s_0))$  where  $x \in \text{bound}(\varphi)$  is a  $\nu$ -variable. Then, if there is a position  $\langle y, s_2 \rangle$  reachable from  $\langle x, s_1 \rangle$  in  $V|_{\sigma^N}$ , where  $y \in \text{bound}(\varphi)$  is a  $\mu$ -variable, then there is no position  $\langle x, s_3 \rangle$  reachable from  $\langle y, s_2 \rangle$  in  $V|_{\sigma^N}$ .*

*Proof.* Suppose there is a play  $\pi$  consistent with  $\sigma^N$  such that we have the following regenerations:  $\langle x, s_1 \rangle$  then  $\langle y, s_2 \rangle$  and then  $\langle x, s_3 \rangle$ , where  $x$  is a  $\nu$ -variable and  $y$  a  $\mu$ -variable. Note that, since  $\varphi \in \Sigma_2^\mu$ , we have that  $y \in \text{free}(\varphi_x)$ , and therefore  $\varphi_x < \varphi_y$ . This implies that in  $\pi$  we must have positions of the form  $\langle\diamond(\beta(x)), s_1'\rangle$  and  $\langle\beta(x), s_1''\rangle$  before  $\langle x, s_1 \rangle$ , and also positions of the form  $\langle\diamond(\beta(x)), s_3'\rangle$  and  $\langle\beta(x), s_3''\rangle$  before  $\langle x, s_3 \rangle$  but after  $\langle y, s_2 \rangle$ . By construction of normalized strategy and by the transitivity of  $\mathcal{T}$  it holds that  $d(\langle\beta(x), s_1''\rangle) = d(\langle\beta(x), s_3''\rangle)$  but also that  $d(\langle\beta(x), s_1''\rangle) = d(\langle\beta(x), s_3''\rangle) = d(\langle y, s_2 \rangle)$ . This implies that  $\text{scc}(\langle y, s_2 \rangle) \neq \emptyset$ . Because  $\sigma^N$  is a winning strategy for Player 0, by Lemma 26 we get the desired contradiction.  $\dashv$

We immediately can restate the previous lemma as the following theorem.

**Theorem 29** *Suppose a finite transitive transition system  $\mathcal{T}$ , a formula  $\varphi \in \Sigma_2^\mu$  such that all  $\nu$ -variables are weakly existential and a normalized winning strategy,  $\sigma^N$ , of Player 0 in  $\mathcal{E}(\varphi, (\mathcal{T}, s))$ . If in a play  $\pi$  consistent with  $\sigma^N$  there is a*

regeneration of a  $\nu$ -variable  $x$  then either there is no more regeneration of a  $\mu$ -variable after the first regeneration of  $x$  or, if there is such a regeneration of a  $\mu$ -variable, then after this position there is no more regeneration of  $x$ .

#### 6.4 Encoding normalized winning strategies

In Definition 5 we define the formulae  $\text{NS}_\varphi^-(X', y)$  and  $\text{NS}_\varphi^+(x, X')$  used to encode the main properties of the normalization of winning strategies of  $\varphi$  given by Theorem 29. Encoding, in this context, will be formalized in the two main Lemmas of the section, Lemmas 31 and 32. The intuition behind these formulae is the following:

- $\text{NS}_\varphi^-(X', y)$  reflects the fact that we are regenerating  $y$  and any  $\nu$ -variable regenerated afterwards will be an element of  $X'$ ,
- $\text{NS}_\varphi^+(x, X')$  reflects the fact that we are regenerating  $x$  and if we regenerate any  $\mu$ -variable then afterwards any  $\nu$ -variable regenerated will be an element of  $X'$ .

In the sequel, in order to ease notation, we write a formula of the form  $\varphi_y^{\text{free}(X)}$  instead of  $(\varphi^{\text{free}(X)})_y$ .

**Definition 5.** Let  $\varphi$  be a  $\Sigma_2^\mu$ -formula. Let  $Y = \{y_1, \dots, y_k\}$  be the set of all  $\mu$ -variables in  $\varphi$  and  $X$  be the set of all  $\nu$ -variables in  $\varphi$ . For all subsets of  $X' \subset X$ , all  $\nu$ -variables  $x$  such that  $x \in X/X'$  and all  $\mu$ -variables  $y$  we define the formulae  $\text{NS}_\varphi^+(x, X')$  and  $\text{NS}_\varphi^-(X', y)$  recursively on the size of  $X'$  such that

$$\text{NS}_\varphi^-(\emptyset, y) \equiv \text{unf}_{\varphi-x}^Y((\varphi^{-X})_y)$$

and, such that

$$\text{NS}_\varphi^+(x, \emptyset) \equiv (\text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)})[y_1/\text{NS}_\varphi^-(\emptyset, y_1), \dots, y_k/\text{NS}_\varphi^-(\emptyset, y_k)].$$

If  $X' = \{x_{i_1}, \dots, x_{i_l}\}$  and  $\overline{X'} = X \setminus X'$ , then

$$\begin{aligned} \text{NS}_\varphi^-(X', y) \equiv & (\text{unf}_{(\varphi-\overline{X'})^{\text{free}(X')}}^Y (\varphi^{-\overline{X'}})_y^{\text{free}(X')})[x_{i_1}/\text{NS}_{\varphi-\overline{X'}}^+(x_{i_1}, X'^{-i_1}), \\ & \vdots \\ & x_{i_l}/\text{NS}_{\varphi-\overline{X'}}^+(x_{i_l}, X'^{-i_l})], \end{aligned}$$

and

$$\begin{aligned} \text{NS}_\varphi^+(x, X') \equiv & (\text{unf}_{\varphi^{\text{free}(Y \cup X')}}^{\overline{X'}} \varphi_x^{\text{free}(Y \cup X')})[y_1/\text{NS}_\varphi^-(X', y_1), \\ & \vdots \\ & y_k/\text{NS}_\varphi^-(X', y_k), \\ & x_{i_1}/\text{NS}_\varphi^+(x_{i_1}, X'^{-i_1}), \\ & \vdots \\ & x_{i_l}/\text{NS}_\varphi^+(x_{i_l}, X'^{-i_l})]. \end{aligned}$$

Note that by construction we have that for every  $\nu$ -variable  $x$ , every  $\mu$ -variable  $y$  and every set of  $\nu$ -variables  $X'$ ,  $\text{free}(\text{NS}_\varphi^+(x, X')), \text{free}(\text{NS}_\varphi^-(X', y)) \subseteq \text{free}(\varphi)$  and  $\text{bound}(\text{NS}_\varphi^+(x, X')), \text{bound}(\text{NS}_\varphi^-(X', y)) \subseteq \text{bound}(\varphi)$ .

**Lemma 30** *Let  $\varphi \in \Sigma_2^\mu$ ,  $y$  be  $\mu$ -variable in  $\varphi$  and  $X'$  be a proper subset of the set of all  $\nu$ -variables. Suppose that  $x_i$  is a  $\nu$ -variable such that  $x_i \notin X'$ . We have that*

$$\text{NS}_\varphi^-(X', y), \text{NS}_\varphi^+(x_i, X') \in \Delta_2^\mu.$$

*Proof.* The proof goes by induction on the size of  $X'$ . If  $X' = \emptyset$  then clearly  $\text{NS}_\varphi^-(X', Y) \in \Sigma_1^\mu$  and, by definition of the formula,  $\text{NS}_\varphi^+(x, X') \in \Delta_2^\mu$ . The induction step follows from the definitions by noting that the class  $\Delta_2^\mu$  is closed under substitution of  $\Delta_2^\mu$  formulae if no new variable is bound.  $\dashv$

**Lemma 31** *Let  $\varphi$  be a  $\Sigma_2^\mu$ -formula and  $X$  be the set of all  $\nu$ -variables in  $\varphi$ . Suppose that all  $x \in X$  are weakly existential. Let  $(\mathcal{T}, s_0)$  be a finite transitive transition system such that there is a normalized winning strategy  $\sigma^N$  in the evaluation game  $\mathcal{E}(\varphi, (\mathcal{T}, s_0))$ . The following holds for every  $X' \subseteq X$  where  $\bar{X}' = X/X'$ :*

1. *If there is a play consistent with  $\sigma^N$  which reaches a position  $\langle y, s \rangle$  ( $y$  a  $\mu$ -variable in  $\varphi$ ) such that on this play before  $\langle y, s \rangle$  there are positions  $\langle \bar{x}, s \rangle$  for all  $\bar{x} \in \bar{X}'$  then it holds that*

$$s \in \|\text{NS}_\varphi^-(X', y)\|_{\mathcal{T}^\varphi}.$$

2. *If there is a play consistent with  $\sigma^N$  which reaches for the first time a position  $\langle x, s \rangle$  ( $x$  a  $\nu$ -variable in  $\varphi$ ) such that on this play before  $\langle x, s \rangle$  there are positions  $\langle \bar{x}, s \rangle$  for all  $\bar{x} \in \bar{X}' \setminus \{x\}$  then it holds that*

$$s \in \|\text{NS}_\varphi^+(x, X')\|_{\mathcal{T}^\varphi}.$$

*Proof.* Let  $Y = \{y_1, \dots, y_k\}$  be the set of all  $\mu$ -variables in  $\varphi$ . We prove the two points simultaneously by induction on the size of  $X'$ . If  $X' = \emptyset$  we have that  $\text{NS}_\varphi^-(\emptyset, y) \equiv \text{unf}_{\varphi-x}^Y((\varphi^{-X})_y)$ . If there is a play consistent with  $\sigma^N$  reaching a position of the form  $\langle y, s \rangle$  whereby for all  $\nu$ -variables there has been a regeneration in this play before, then, since  $\sigma^N$  is a normalized strategy, by Theorem 29 there can not be any regeneration of a  $\nu$ -variable after  $\langle y, s \rangle$ . Therefore  $\sigma^N$  determines a winning strategy in

$$\mathcal{E}(\text{unf}_{\varphi-x}^Y((\varphi^{-X})_y), (\mathcal{T}, s))$$

and with Theorem 7 we get the induction base for part 1. For part 2 remember that

$$\text{NS}_\varphi^+(x, \emptyset) \equiv (\text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)})[y_1/\text{NS}_\varphi^-(\emptyset, y_1), \dots, y_k/\text{NS}_\varphi^-(\emptyset, y_k)].$$

Suppose that there is a play consistent with  $\sigma^N$  which reaches for the first time a position  $\langle x, s \rangle$  ( $x \in X$ ) such that on this play before  $\langle x, s \rangle$  there are positions

$\langle \bar{x}, s \rangle$  for all  $\bar{x} \in X \setminus \{x\}$ . Then, since  $\sigma^N$  is a normalized strategy, by Theorem 29 for every play extending this position which is compatible with  $\sigma^N$ , either there are only regenerations of  $\nu$ -variables, or, if there is a regeneration of a  $\langle y, s \rangle$ , then after this regeneration there is no more regeneration of a  $\nu$ -variable. Therefore  $\sigma^N$  determines a winning strategy in

$$\mathcal{E}((\text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)})[y_1/\text{NS}_{\varphi}^-(\emptyset, y_1), \dots, y_k/\text{NS}_{\varphi}^-(\emptyset, y_k)], (\mathcal{T}, s))$$

and with Theorem 7 we get the induction base for part 2.

For the induction step of part 1, let  $X' = \{x_{i_1}, \dots, x_{i_l}\}$  and let  $\langle y, s \rangle$  be a position of a play consistent with  $\sigma^N$  such that all  $\bar{x} \in \bar{X}'$  have been regenerated before. Then, by Theorem 29 for all  $\nu$ -variables  $x_i$  regenerated afterwards in the play we have  $x_i \in X'$ . By construction for such a position  $\langle x_i, s_i \rangle$  we will have that all  $\nu$ -variables in  $\bar{X}'$  are regenerated before this position. Define  $X'^{-i} = X' \setminus \{x_i\}$ . It can easily be seen that  $\langle x_i, s_i \rangle$  satisfy the condition of part 2 and, since  $x_i \in X'$ , that  $X'^{-i} \subsetneq X'$ . Therefore, we can apply induction hypothesis of part 2 and get

$$s_i \in \|\text{NS}^+(x_i, X'^{-i})\|_{\mathcal{T}}.$$

Recapitulating, we have that for all plays consistent with  $\sigma^N$  starting from  $\langle y, s \rangle$  if a  $\nu$ -variable  $x_i$  is regenerated by a position  $\langle x_i, s_i \rangle$  then we have  $s_i \in \|\text{NS}^+(x_i, X'^{-i})\|_{\mathcal{T}}$  and otherwise we have only regenerations of  $\mu$ -variables. But by Lemmas 3.1 and 8 this means that  $\sigma^N$  gives us a winning strategy in the evaluation game

$$\mathcal{E}(\gamma, (\mathcal{T}, s))$$

where

$$\begin{aligned} \gamma \equiv \text{unf}_{(\varphi^{-\bar{X}'})_{\text{free}(X')}}^Y ((\varphi^{-\bar{X}'})_y^{\text{free}(X')}) [ & x_{i_1}/\text{NS}_{\varphi^{-\bar{X}'}}^+(x_{i_1}, X'^{-i_1}), \\ & \vdots \\ & x_{i_l}/\text{NS}_{\varphi^{-\bar{X}'}}^+(x_{i_l}, X'^{-i_l})]. \end{aligned}$$

By noting that  $\gamma \equiv \text{NS}_{\varphi}^-(X', y)$  and using Theorem 7 we finish the induction step for part 1.

For the induction step of part 2 let  $\langle x, s \rangle$  be a position of a play consistent with  $\sigma^N$  such that all  $\bar{x} \in \bar{X}'$  have been regenerated before. There are only three disjoint classes of winning plays (consistent with  $\sigma^N$ ) extending the position  $\langle x, s \rangle$  and they are obtained by considering all possible regenerations of bound variables after this position:

1. The class of plays in which afterwards we regenerate a  $x_i \in X'$  in a position  $\langle x_i, s_i \rangle$ , and before this position there was no regeneration of a  $\mu$ -variable. In this case we can apply the induction hypothesis for part 2 to the set  $X'^{-i}$  and get

$$s_i \in \|\text{NS}_{\varphi}^+(x_i, X'^{-i})\|_{\mathcal{T}}.$$

2. The class of plays in which afterwards we regenerate a  $\mu$ -variable  $y$  in a position  $\langle y, s_y \rangle$ , and before this position there was no regeneration of a  $x_i \in X'$ . In this case, we can apply part 1, where the induction step is already done, and get

$$s_y \in \|\text{NS}_\varphi^-(X', y)\|_{\mathcal{T}}.$$

3. The class of plays in which there is no regeneration of  $z \in X' \cup Y$ , but there are eventually only regenerations of  $x_i \in \overline{X'}$ . Because these plays are consistent with  $\sigma^N$ , they are winning. Therefore, they are winning in the evaluation game  $\mathcal{E}((\text{unf}_{\varphi^-(Y \cup X')}^{\overline{X'}} \varphi_x^-(Y \cup X')), (\mathcal{T}, s))$ , too.

By Lemmas 3.1 and 8 we have that

$$s \in \left\| \left( \text{unf}_{\varphi^{\text{free}(Y \cup X')}}^{\overline{X'}} \varphi_x^{\text{free}(Y \cup X')} \right) \left[ \begin{array}{l} y_1 / \text{NS}_\varphi^-(X', y_1), \\ \vdots \\ y_k / \text{NS}_\varphi^-(X', y_k), \\ x_{i_1} / \text{NS}_\varphi^+(x_{i_1}, X'^{-i_1}), \\ \vdots \\ x_{i_l} / \text{NS}_\varphi^+(x_{i_l}, X'^{-i_l}) \end{array} \right] \right\|_{\mathcal{T}}.$$

and this ends the induction step of part 2 and the proof.  $\dashv$

**Lemma 32** *Let  $\varphi$  be a  $\Sigma_2^\mu$ -formula and  $X$  be the set of all  $\nu$ -variables in  $\varphi$ . Suppose that all  $\nu$ -variables are weakly existential. Then, for every finite transitive transition system  $\mathcal{T}$  and for every  $X' \subseteq X$  it holds that*

1. For every  $y \in Y$  we have

$$\|\text{NS}_\varphi^-(X', y)\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_y\|_{\mathcal{T}^\varphi}, \text{ and}$$

2. for every  $x \in \overline{X'} =: X/X'$  we have

$$\|\text{NS}_\varphi^+(x, X')\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}.$$

*Proof.* Let  $Y = \{y_1, \dots, y_k\}$  be the set of all  $\mu$ -variables. We prove the two points simultaneously by induction on the size of  $X'$ . Suppose  $X'$  is empty. Then we have that  $\text{NS}_\varphi^-(\emptyset, y) \equiv \text{unf}_{\varphi^{-X}}^Y((\varphi^{-X})_y)$  and by Lemma 22.3 we obtain

$$\|\text{unf}_{\varphi^{-X}}^Y((\varphi^{-X})_y)\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_y\|_{\mathcal{T}^\varphi}.$$

Therefore we complete the base case of the induction for part 1. For part 2 remember that

$$\text{NS}_\varphi^+(x, \emptyset) \equiv (\text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)})[y_1 / \text{NS}_\varphi^-(\emptyset, y_1), \dots, y_k / \text{NS}_\varphi^-(\emptyset, y_k)].$$

Thus, by the induction base of part 1 and by Lemma 3.4, we have that

$$\begin{aligned} & \left\| \left( \text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)} \right) [y_1 / \text{NS}_\varphi^-(\emptyset, y_1), \dots, y_k / \text{NS}_\varphi^-(\emptyset, y_k)] \right\|_{\mathcal{T}^\varphi} \\ & \subseteq \\ & \left\| \left( \text{unf}_{\varphi^{\text{free}(Y)}}^X \varphi_x^{\text{free}(Y)} \right) [y_1 / \varphi_{y_1}, \dots, y_m / \varphi_{y_m}] \right\|_{\mathcal{T}^\varphi}. \end{aligned}$$

But because in  $\mathcal{T}^\varphi$  we have that  $\lambda(y) = \|\varphi_y\|_{\mathcal{T}^\varphi}$  and by applying Lemma 3.1 and Lemma 21.2, it holds that

$$\|(\mathbf{unf}_{\varphi^{\text{free}(Y)} \varphi_x^{\text{free}(Y)}}^X)[y_1/\varphi_{y_1}, \dots, y_m/\varphi_{y_m}]\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}.$$

Therefore

$$\|(\mathbf{unf}_{\varphi^{\text{free}(Y)} \varphi_x^{\text{free}(Y)}}^X)[y_1/\mathbf{NS}_\varphi^-(\emptyset, y_1), \dots, y_k/\mathbf{NS}_\varphi^-(\emptyset, y_k)]\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}.$$

This ends the induction base for both parts 1 and 2.

Let  $X' = \{x_{i_1}, \dots, x_{i_l}\}$ . For the induction step of part 1, remember that

$$\begin{aligned} \mathbf{NS}_\varphi^-(X', y) \equiv \mathbf{unf}_{(\varphi^{-\overline{X'}})^{\text{free}(X')}}^Y((\varphi^{-\overline{X'}})^{\text{free}(X')})_y [x_{i_1}/\mathbf{NS}_{\varphi^{-\overline{X'}}}^+(x_{i_1}, X'^{-i_1}), \\ \vdots \\ x_{i_l}/\mathbf{NS}_{\varphi^{-\overline{X'}}}^+(x_{i_l}, X'^{-i_l})]. \end{aligned}$$

By induction hypothesis, by Lemma 3.4 and because in  $\mathcal{T}^{\varphi^{-\overline{X'}}$  the evaluation of a variable  $x_{i_j} \in X'$  is equal to  $\|(\varphi^{-\overline{X'}})_{x_{i_j}}\|_{\mathcal{T}^{\varphi^{-\overline{X'}}$ , we obtain

$$\begin{aligned} \|\mathbf{NS}_\varphi^-(X', y)\|_{\mathcal{T}^\varphi} &= \|\mathbf{unf}_{(\varphi^{-\overline{X'}})^{\text{free}(X')}}^Y((\varphi^{-\overline{X'}})^{\text{free}(X')})_y \\ &\quad [x_{i_1}/\mathbf{NS}_{\varphi^{-\overline{X'}}}^+(x_{i_1}, X'^{-i_1}), \\ &\quad \vdots \\ &\quad x_{i_l}/\mathbf{NS}_{\varphi^{-\overline{X'}}}^+(x_{i_l}, X'^{-i_l})]\|_{\mathcal{T}} \\ &\subseteq \|\mathbf{unf}_{(\varphi^{-\overline{X'}})^{\text{free}(X')}}^Y((\varphi^{-\overline{X'}})^{\text{free}(X')})_y\|_{\mathcal{T}^{\varphi^{-\overline{X'}}}}. \end{aligned}$$

With Lemma 21.2 we obtain

$$\|\mathbf{unf}_{(\varphi^{-\overline{X'}})^{\text{free}(X')}}^Y((\varphi^{-\overline{X'}})^{\text{free}(X')})_y\|_{\mathcal{T}^{\varphi^{-\overline{X'}}}} = \|(\varphi^{-\overline{X'}})_y\|_{\mathcal{T}^{\varphi^{-\overline{X'}}}}.$$

Finally, because by Lemma 22.4 it holds that  $\|(\varphi^{-\overline{X'}})_y\|_{\mathcal{T}^{\varphi^{-\overline{X'}}}} \subseteq \|\varphi_y\|_{\mathcal{T}^\varphi}$  we get

$$\|\mathbf{NS}_\varphi^-(X', y)\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_y\|_{\mathcal{T}^\varphi}.$$

For the induction step of part 2 if  $\overline{X} = X \setminus X'$ , then by induction hypothesis and by part 1 we have for every finite transitive transition system  $\mathcal{T}$

$$\begin{aligned} \|\mathbf{NS}_\varphi^-(X', y_1)\|_{\mathcal{T}^\varphi} &\subseteq \|\varphi_{y_1}\|_{\mathcal{T}^\varphi}, \\ &\vdots \\ \|\mathbf{NS}_\varphi^-(X', y_k)\|_{\mathcal{T}^\varphi} &\subseteq \|\varphi_{y_k}\|_{\mathcal{T}^\varphi}, \\ \|\mathbf{NS}_\varphi^+(x_{i_1}, X'^{-i_1})\|_{\mathcal{T}^\varphi} &\subseteq \|\varphi_{x_{i_1}}\|_{\mathcal{T}^\varphi}, \\ &\vdots \\ \|\mathbf{NS}_\varphi^+(x_{i_l}, X'^{-i_l})\|_{\mathcal{T}^\varphi} &\subseteq \|\varphi_{x_{i_l}}\|_{\mathcal{T}^\varphi}. \end{aligned}$$

Therefore, by Lemmas 3.4 and 3.1, and because in  $\mathcal{T}^\varphi$  we have that  $\lambda(z) = \|\varphi_z\|_{\mathcal{T}^\varphi}$ , for every  $z \in \text{bound}(\varphi)$ , we get

$$\begin{aligned} & \|(\text{unf}_{\varphi^{\overline{X'}}}^{\text{free}(Y \cup X')}(\varphi_x^{\text{free}(Y \cup X')})) [ \\ & \quad y_1/\text{NS}_{\varphi}^-(X', y_1), \\ & \quad \vdots \\ & \quad y_k/\text{NS}_{\varphi}^-(X', y_k) \\ & \quad x_{i_1}/\text{NS}_{\varphi}^+(x_{i_1}, X'^{-i_1}), \\ & \quad \vdots \\ & \quad x_{i_l}/\text{NS}_{\varphi}^+(x_{i_l}, X'^{-i_l}) ] \|_{\mathcal{T}^\varphi} \\ & \subseteq \| \text{unf}_{\varphi^{\overline{X'}}}^{\text{free}(Y \cup X')}(\varphi_x^{\text{free}(Y \cup X')}) \|_{\mathcal{T}^\varphi}. \end{aligned}$$

We can apply Lemma 21.2 and obtain

$$\|(\text{unf}_{\varphi^{\overline{X'}}}^{\text{free}(Y \cup X')}(\varphi_x^{\text{free}(Y \cup X')}))\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}.$$

Because this implies that

$$\|\text{NS}_{\varphi}^+(x, X')\|_{\mathcal{T}^\varphi} \subseteq \|\varphi_x\|_{\mathcal{T}^\varphi}$$

this ends the induction step of part 2 and the proof of the Lemma.  $\dashv$

## 6.5 The collapse over transitive models

Everything now is ready to prove the collapse of the  $\mu$ -hierarchy over finite transitive transition systems.

**Definition 6.** For the formula  $\varphi \in \Sigma_2^\mu$  such that  $X = \{x_1, \dots, x_m\}$  is the set of all  $\nu$ -variables in  $\varphi$ . We define a new formula  $\rho(\varphi) \in \Delta_2^\mu$  such that

$$\rho(\varphi) \equiv \varphi^{\text{free}(X)}[x_1/\text{NS}_{\varphi}^+(x_1, X^{-1}), \dots, x_m/\text{NS}_{\varphi}^+(x_m, X^{-m})].$$

*Remark 3.* By Lemma 30 it can easily be seen that  $\rho(\varphi)$  is indeed a  $\Delta_2^\mu$ -formula.

**Theorem 33** For all  $\varphi \in \Sigma_2^\mu$  and all finite transitive transition systems  $\mathcal{T}$  we have that

$$\|\varphi\|_{\mathcal{T}} = \|\rho(\tau(\varphi))\|_{\mathcal{T}}.$$

*Proof.* First, we observe that  $\tau(\varphi) \in \Sigma_2^\mu$  and that by Corollary 25 we have that  $\|\varphi\|_{\mathcal{T}} = \|\tau(\varphi)\|_{\mathcal{T}}$ . Thus, we can assume that each  $\nu$ -variable in  $\varphi \in \Sigma_2^\mu$  is weakly existential and any  $\mu$ -variable weakly universal. If  $X = \{x_1, \dots, x_m\}$  is the set of all  $\nu$ -variables in  $\varphi$ , by definition of  $\rho$  we have to prove that

$$\|\varphi\|_{\mathcal{T}} = \|\varphi^{\text{free}(X)}[x_1/\text{NS}_{\varphi}^+(x_1, X^{-1}), \dots, x_m/\text{NS}_{\varphi}^+(x_m, X^{-m})]\|_{\mathcal{T}}.$$

“ $\supseteq$ ”: First note that because for every  $x_i \in X$  it holds that  $\|\text{NS}_{\varphi}^+(x_i, X^{-i})\|_{\mathcal{T}} = \|\text{NS}_{\varphi}^+(x_i, X^{-i})\|_{\mathcal{T}^\varphi}$ , the models  $\mathcal{T}[x_1 \mapsto \|\text{NS}_{\varphi}^+(x_1, X^{-1})\|_{\mathcal{T}}, \dots, x_m \mapsto \|\text{NS}_{\varphi}^+(x_m, X^{-m})\|_{\mathcal{T}}]$

and  $\mathcal{T}^\varphi[x_1 \mapsto \|\mathbf{NS}_\varphi^+(x_1, X^{-1})\|_{\mathcal{T}^\varphi}, \dots, x_m \mapsto \|\mathbf{NS}_\varphi^+(x_m, X^{-m})\|_{\mathcal{T}^\varphi}]$  agree on the free variables of  $\varphi^{\text{free}(X)}$ . Therefore we have

$$\begin{aligned} & \|\varphi^{\text{free}(X)}[x_1/\mathbf{NS}_\varphi^+(x_1, X^{-1}), \dots, x_m/\mathbf{NS}_\varphi^+(x_m, X^{-m})]\|_{\mathcal{T}} = \\ & \|\varphi^{\text{free}(X)}[x_1/\mathbf{NS}_\varphi^+(x_1, X^{-1}), \dots, x_m/\mathbf{NS}_\varphi^+(x_m, X^{-m})]\|_{\mathcal{T}^\varphi} \end{aligned}$$

With Lemma 32 and, because all  $\nu$ -variables appear positively in  $\varphi$ , by applying Lemma 3.4 we get that

$$\begin{aligned} & \|\varphi^{\text{free}(X)}[x_1/\mathbf{NS}_\varphi^+(x_1, X^{-1}), \dots, x_m/\mathbf{NS}_\varphi^+(x_m, X^{-m})]\|_{\mathcal{T}^\varphi} \\ & \quad \subseteq \\ & \|\varphi^{\text{free}(X)}[x_1/\varphi_{x_1}, \dots, x_m/\varphi_{x_m}]\|_{\mathcal{T}^\varphi} \end{aligned}$$

By Lemma 3.4 and because in  $\mathcal{T}^\varphi$  we have that  $\lambda(x_i) = \|\varphi_{x_i}\|_{\mathcal{T}^\varphi}$ , we obtain

$$\|\varphi^{\text{free}(X)}[x_1/\varphi_{x_1}, \dots, x_m/\varphi_{x_m}]\|_{\mathcal{T}^\varphi} \subseteq \|\varphi^{\text{free}(X)}\|_{\mathcal{T}^\varphi}.$$

Since by Lemma 21.2 we have that  $\|\varphi^{\text{free}(X)}\|_{\mathcal{T}^\varphi} = \|\varphi\|_{\mathcal{T}}$  we get this inclusion.

“ $\subseteq$ ”: Let  $s \in \|\varphi\|_{\mathcal{T}}$ . By Theorem 7 there is a winning strategy in  $\mathcal{E}(\varphi, (\mathcal{T}, s))$  and by Theorem 29 it can be assumed to be normalized. Let  $\pi$  be any play consistent with the strategy starting from  $\langle \varphi, s \rangle$ . We have that if there is a (first) regeneration of a  $\nu$ -variable  $x_i$  in a position  $\langle x_i, s_i \rangle$  then by Lemma 31 we have that

$$s_i \in \|\mathbf{NS}^+(x_i, X^{-i})\|_{\mathcal{T}}$$

where  $X$  is the set of all  $\nu$ -variables in  $\varphi$ . Therefore, there is a winning strategy for Player 0 in

$$\mathcal{E}(\varphi^{\text{free}(X)}, (\mathcal{T}[x_1 \mapsto \|\mathbf{NS}^+(x_1, X^{-1})\|_{\mathcal{T}}, \dots, x_n \mapsto \|\mathbf{NS}^+(x_n, X^{-n})\|_{\mathcal{T}}], s))$$

By Theorem 7 we have that

$$s \in \|\varphi^{\text{free}(X)}\|_{\mathcal{T}[x_1 \mapsto \|\mathbf{NS}^+(x_1, X^{-1})\|_{\mathcal{T}}, \dots, x_n \mapsto \|\mathbf{NS}^+(x_n, X^{-n})\|_{\mathcal{T}}]}$$

and with Lemma 3.1 we complete the proof.  $\dashv$

**Corollary 34** *The modal  $\mu$ -calculus hierarchy on finite transitive systems collapses to  $\Delta_2^\mu$ .*

*Proof.* By Theorem 33 it holds that  $\Sigma_2^{\mu\mathbb{T}^{tf}} = \Delta_2^{\mu\mathbb{T}^{tf}}$ . By duality,  $\Pi_2^{\mu\mathbb{T}^{tf}} = \Delta_2^{\mu\mathbb{T}^{tf}}$ . By this fact it is therefore very easy to verify inductively that for every  $n > 0$ ,  $\Sigma_{2+n}^{\mu\mathbb{T}^{tf}} = \Pi_{2+n}^{\mu\mathbb{T}^{tf}} = \Delta_2^{\mu\mathbb{T}^{tf}}$ .  $\dashv$

**Corollary 35** *The modal  $\mu$ -calculus hierarchy on transitive systems collapses to  $\Delta_2^\mu$ .*

*Proof.* Suppose that the hierarchy does not collapse. Therefore, there is a formula  $\varphi$  such that for all formula  $\psi \in \Delta_2^\mu$  there is a transitive system  $\mathcal{T}$  such that  $\mathcal{T}, s_0 \models \neg(\varphi \leftrightarrow \psi)$ . By Theorem 16, there is a finite transitive model  $\mathcal{T}^f$  such that  $\mathcal{T}^f, s_i^f \models \neg(\varphi \leftrightarrow \psi)$ . But this cannot be the case by Corollary 34.  $\dashv$



We end with the definition of a syntactical translation from  $\mathcal{L}_\mu$  to  $\Delta_2^\mu$  preserving equivalence on transitive transition systems.

**Definition 7.**  $R : \mathcal{L}_\mu \rightarrow \Delta_2^\mu$  is defined as

- $R(p) = p$  and  $R(\neg p) = \neg p$
- $R(\perp) = \perp$  and  $R(\top) = \top$
- $R(\alpha \circ \beta) = R(\alpha) \circ R(\beta)$ , where  $\circ \in \{\wedge, \vee\}$
- $R(\Delta \beta) = \Delta R(\beta)$ , where  $\Delta \in \{\Box, \Diamond\}$
- $R(\mu x.\varphi) = \text{wn}(\rho(\tau(\text{wn}(\mu x.(R(\varphi))))))$
- $R(\nu x.\varphi) = \neg(R(\mu x.\neg\varphi[x/\neg x]))$

**Lemma 36** For all  $\mu$ -formula  $\varphi$  we have that

1.  $R(\varphi)$  is well-defined, and
2.  $R(\varphi) \in \Delta_2^\mu$ .

*Proof.* We prove both parts simultaneously by induction on the structure of  $\varphi$ . The induction cases for boolean and modal connectives are trivial. If  $\varphi$  is of the form  $\mu x.\alpha$  we have that  $R(\mu x.\alpha) = \text{wn}(\rho(\tau(\text{wn}(\mu x.(R(\alpha))))))$ . Because  $\tau$  is a well-defined syntactical transformation, and neither  $\text{wn}$  nor  $\tau$  increase the alternation depth of a formula, the application of  $\rho$  in the clause of  $R(\mu x.\alpha)$  is well-defined by induction hypothesis. Thus,  $R(\varphi)$  terminates and therefore it is well-defined too. The fact that  $R(\mu x.\alpha) \in \Delta_2^\mu$  follows by induction hypothesis, by the fact that, by Remark 3, for all  $\Sigma_2^\mu$ -formulae  $\psi$  we have that  $\rho(\psi) \in \Delta_2^\mu$ , and because we know that  $\tau$  and  $\text{wn}$  do not increase the alternation depth. If  $\varphi$  is of the form  $\nu x.\alpha$ , on one hand  $R(\nu x.\alpha)$  is well-defined because the clause for this form is defined via a reducing case  $R(\mu x.\neg\varphi[x/\neg x])$ , and, on the other hand  $R(\varphi) \in \Delta_2^\mu$  because  $\Delta_2^\mu$  is closed under negation.  $\dashv$

**Theorem 37** For all  $\varphi \in \mathcal{L}_\mu$  and all finite transitive transition systems  $\mathcal{T}$  we have that

$$\|\varphi\|_{\mathcal{T}} = \|R(\varphi)\|_{\mathcal{T}}.$$

*Proof.* We prove the equivalence by induction on  $\text{rank}(\varphi)$  simultaneously for all finite transitive transition systems  $\mathcal{T}$ . The induction cases for boolean and modal connectives are trivial. If  $\varphi$  is of the form  $\mu x.\alpha$  we have that

$$\begin{aligned} \|R(\mu x.\alpha)\|_{\mathcal{T}} &= \|\text{wn}(\rho(\tau(\text{wn}(\mu x.R(\alpha)))))\|_{\mathcal{T}} && \text{by definition of } R \\ &= \|\text{wn}(\mu x.R(\alpha))\|_{\mathcal{T}} && \tau(\text{wn}(\mu x.R(\alpha))) \in \Sigma_2^\mu, \text{ Lemma 4} \\ &&& \text{and Theorem 33} \\ &= \|\mu x.\alpha\|_{\mathcal{T}} && \text{by Lemma 4 and induction hypothesis} \end{aligned}$$

If  $\varphi$  is of the form  $\nu x.\alpha$  we do a similar induction step like above by using the equivalence  $\|\nu x.\alpha\|_{\mathcal{T}} = \|\neg\mu x.\neg\alpha[x/\neg x]\|_{\mathcal{T}}$ .  $\dashv$

We conclude by verifying that the syntactical translation  $R$  is also an explicit syntactical translation of all modal  $\mu$ -formulae to the alternation free fragment preserving denotation in *every* transitive transition systems. The proof goes with similar argument as in Corollary 35 and it is left to the reader.

**Theorem 38** For all  $\varphi \in \mathcal{L}_\mu$  and all transitive transition systems  $\mathcal{T}$  we have that

$$\|\varphi\|_{\mathcal{T}} = \|R(\varphi)\|_{\mathcal{T}}.$$

*Remark 4.* Note, that due to the example of Visser in [Vis96] mentioned in the introduction the alternation-free fragment is also the optimal bound if restrict ourselves to transition systems which are transitive *and* reflexive.

*Example 5.* Let's have a look at our example from Section 2. In the case of “always eventually”, we have that

$$\|\nu x.(\mu y.(p \vee \diamond y)) \wedge \square x\|^{\mathbb{T}^t} = \|(p \vee \diamond p) \wedge \square(p \vee \diamond p)\|^{\mathbb{T}^t}.$$

For “infinitely often”, it holds that

$$\|\nu x.\mu y.((p \vee \diamond y)) \wedge \diamond x\|^{\mathbb{T}^t} = \|\nu x.(p \wedge \diamond x)\|^{\mathbb{T}^t}.$$

But, because from footnote 4 of the introduction we know that  $\nu x.(p \wedge \diamond x)$  cannot be reduced to any purely modal formula, contrary to the transitive *and* symmetric case, over transitive transition systems “infinitely often” cannot be expressed by a  $\Delta_1^\mu$  formula.

## 7 Strictness of the hierarchy for reflexive transition systems

In this section we prove the strictness of the modal  $\mu$ -calculus hierarchy on reflexive transition systems. In doing this, we follow the argumentation of the proof of the strictness of the hierarchy on all binary transition systems presented in [Alb02]. First, we adapt the game transition system such that it is reflexive.

Let  $\mathcal{E}(\varphi, (\mathcal{T}, s))$  be a parity game with priority function  $\Omega$  and with corresponding pointed game transition system  $\mathcal{T}(\mathcal{E}(\varphi, (\mathcal{T}, s)))$ . We extend the edge relation  $E$  of the parity game to its reflexive closure  $E^r = E \cup \{(s, s); s \in V_0 \cup V_1\}$ , and change our priority function  $\Omega$  to  $\Omega^r$  such that for all vertices  $\langle \psi, s \rangle$  where  $\psi \equiv \eta x.\delta$  ( $\eta \in \{\mu, \nu\}$ ) we have

$$\Omega^r(\langle \psi, s \rangle) = \Omega(\langle \psi, s \rangle) + 2$$

and such that for all other vertices we define:

– if  $\min \Omega$  is even

$$\Omega^r(\langle \psi, s \rangle) = \begin{cases} 0 & \text{if } \langle \psi, s \rangle \in V_1 \\ 1 & \text{if } \langle \psi, s \rangle \in V_0. \end{cases}$$

– if  $\min \Omega$  is odd

$$\Omega^r(\langle \psi, s \rangle) = \begin{cases} 2 & \text{if } \langle \psi, s \rangle \in V_1 \\ 1 & \text{if } \langle \psi, s \rangle \in V_0. \end{cases}$$

The new resulting “reflexive” parity game is denoted as  $\mathcal{E}^r(\varphi, (\mathcal{T}, s))$ . The following Lemma can be proved by unwinding the definition of winning strategy.

**Lemma 39** *Player 0 has a winning strategy for  $\mathcal{E}^r(\varphi, (\mathcal{T}, s))$  iff Player 0 has a winning strategy for  $\mathcal{E}(\varphi, (\mathcal{T}, s))$ .*

Given a “reflexive” parity game  $\mathcal{E}^r(\varphi, (\mathcal{T}, s))$  the *pointed game transition system*  $\mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s)))$  is defined analogously as above. Obviously, the pointed game transition system  $\mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s)))$  is reflexive. We have that

**Proposition 40** *Let  $(\mathcal{T}, s)$  be an arbitrary pointed transition system. For all  $\varphi \in \Pi_n^\mu$  we have that*

$$\mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s))) \in \|W_{\Pi_{n+2}^\mu}\| \quad \text{if and only if} \quad \mathcal{T}(\mathcal{E}(\varphi, (\mathcal{T}, s))) \in \|W_{\Pi_n^\mu}\|$$

and dually for  $\varphi \in \Sigma_n^\mu$ .

*Proof.* This follows directly by the definition of the “reflexive” parity game  $\mathcal{E}^r(\varphi, (\mathcal{T}, s))$  and by applying Proposition 9 to Lemma 39.  $\dashv$

**Corollary 41** *Let  $(\mathcal{T}, s)$  be an arbitrary pointed transition system. For all  $\varphi \in \Pi_n^\mu$  we have that:*

$$\mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s))) \in \|W_{\Pi_{n+2}^\mu}\| \quad \text{if and only if} \quad (\mathcal{T}, s) \in \|\varphi\|.$$

and dually for  $\varphi \in \Sigma_n^\mu$ .

*Proof.* By Proposition 40 and Corollary 11 we obtain our result.  $\dashv$

For all formulae  $\varphi$  we define a function  $f_\varphi$  (functional class) mapping a pointed transition system  $(\mathcal{T}, s)$  to a reflexive transition system  $f_\varphi(\mathcal{T}, s)$  such that

$$f_\varphi(\mathcal{T}, s) := \mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s))).$$

The proof of the next Lemma follows similar arguments as the proof of the same result for arbitrary transition systems proved by one of the authors in [Alb02].

**Lemma 42** *For all formulae  $\psi \in \Sigma_n^\mu$  (resp.  $\Pi_n^\mu$ ),  $n \in \mathbb{N}$ , there is an equivalent formula  $\varphi \in \Sigma_n^\mu$  (resp.  $\Pi_n^\mu$ ) such that the function  $f_\varphi$  has a fixpoint in  $\mathbb{T}^r$ , that is, a pointed reflexive transition system  $(\mathcal{T}^F, s^F)$  such that*

$$f_\varphi(\mathcal{T}^F, s^F) = (\mathcal{T}^F, s^F).$$

**Theorem 43** *For all natural numbers  $n \in \mathbb{N} \setminus \{0\}$  we have that*

$$\Sigma_n^{\mathbb{T}^r} \subsetneq \Sigma_{n+1}^{\mathbb{T}^r} \quad \text{and} \quad \Pi_n^{\mathbb{T}^r} \subsetneq \Pi_{n+1}^{\mathbb{T}^r}.$$

*Proof.* We proof the contrapositive. Assume that we have

$$\Sigma_{n+1}^{\mathbb{T}^r} \subseteq \Sigma_n^{\mathbb{T}^r} \quad \text{or} \quad \Pi_{n+1}^{\mathbb{T}^r} \subseteq \Pi_n^{\mathbb{T}^r}.$$

Without restriction of generality, assume  $\Sigma_{n+1}^{\mathbb{T}^r} \subseteq \Sigma_n^{\mathbb{T}^r}$ . Then, if  $\|\varphi\| \in \Pi_{n+1}^\mu$  we have  $\|\neg\varphi\| \in \Sigma_{n+1}^\mu$  and by assumption  $\|\neg\varphi\| \in \Sigma_n^\mu$  and therefore  $\|\varphi\| \in \Pi_n^\mu$ . Therefore, assuming the contrapositive leads to

$$\Sigma_{n+1}^{\mathbb{T}^r} \subseteq \Sigma_n^{\mathbb{T}^r} \quad \text{and} \quad \Pi_{n+1}^{\mathbb{T}^r} \subseteq \Pi_n^{\mathbb{T}^r}.$$

Since from  $\Sigma_{n+1}^{\mathbb{T}^r} \subseteq \Sigma_n^{\mathbb{T}^r}$ , by definition, it can be inferred that  $\Pi_n^{\mathbb{T}^r} \subseteq \Sigma_n^{\mathbb{T}^r}$ , and from  $\Pi_{n+1}^{\mathbb{T}^r} \subseteq \Pi_n^{\mathbb{T}^r}$ , by definition, it can be inferred that  $\Sigma_n^{\mathbb{T}^r} \subseteq \Pi_n^{\mathbb{T}^r}$ , by assuming the contrapositive we get that  $\Pi_{n+1}^{\mathbb{T}^r} = \Pi_n^{\mathbb{T}^r} = \Sigma_{n+1}^{\mathbb{T}^r} = \Sigma_n^{\mathbb{T}^r}$  and, obviously, we then have for all  $k \in \mathbb{N}$  that

$$\Pi_{n+k}^{\mathbb{T}^r} = \Pi_n^{\mathbb{T}^r} = \Sigma_{n+k}^{\mathbb{T}^r} = \Sigma_n^{\mathbb{T}^r}. \quad (4)$$

Since  $W_{\Sigma_{n+2}^\mu} \in \Sigma_{n+2}^\mu$  we have that  $\neg W_{\Sigma_{n+2}^\mu} \in \Pi_{n+2}^\mu$  and with equation 4 we get

$$\|\neg W_{\Sigma_{n+2}^\mu}\|^{\mathbb{T}^r} \in \Sigma_n^{\mathbb{T}^r}.$$

By Lemma 42 there is a formula  $\varphi \in \Sigma_n^\mu$  equivalent to  $\neg W_{\Sigma_{n+2}^\mu}$  and a pointed transition system  $(\mathcal{T}^F, s^F)$  such that

$$(\mathcal{T}^F, s^F) = f_\varphi(\mathcal{T}^F, s^F).$$

Since  $f_\varphi(\mathcal{T}, s)$  is defined as  $\mathcal{T}(\mathcal{E}^r(\varphi, (\mathcal{T}, s)))$ , by Corollary 41, for all pointed transition systems  $(\mathcal{T}, s)$  we have that  $f_\varphi(\mathcal{T}, s) \in \|W_{\Sigma_{n+2}^\mu}\|$  if and only if  $(\mathcal{T}, s) \in \|\varphi\|$ . Since  $\varphi$  is equivalent to  $\neg W_{\Sigma_{n+2}^\mu}$  we get that

$$(\mathcal{T}^F, s^F) \in \|\neg W_{\Sigma_{n+2}^\mu}\| \quad \text{iff} \quad (\mathcal{T}^F, s^F) \in \|W_{\Sigma_{n+2}^\mu}\|$$

which is a contradiction.  $\dashv$

**Theorem 44** 1. *The modal  $\mu$ -calculus hierarchy is strict over reflexive transition systems.*

2. *The modal  $\mu$ -calculus hierarchy is strict over finite reflexive transition systems.*

*Proof.* Part 1 is a corollary of Theorem 43. For Part 2, let  $\|\varphi\| \in \Sigma_n^{\mathbb{T}^r} \setminus \Pi_n^{\mathbb{T}^r}$ . Then, by Part 1 we know that for every  $\psi \in \Sigma_{n-1}^\mu$  it holds that  $\neg(\varphi \leftrightarrow \psi)$  has a reflexive model. By Theorem 14, this model can be finite. Hence  $\varphi \in \Sigma_n^\mu$  is not equivalent to any  $\Sigma_{n-1}^\mu$  formula on finite reflexive transition systems.  $\dashv$

## Acknowledgements

We thank Igor Walukiewicz, and particularly Giovanna D'Agostino and Giacomo Lenzi for various comments on a preliminary version of this paper.

## References

- [Alb02] ALBERUCCI L.: Strictness of the Modal  $\mu$ -Calculus Hierarchy. in [GTW02], pp. 185–201
- [Alb08] ALBERUCCI L.: A Syntactical Treatment of Simultaneous Fixpoints in the Modal  $\mu$ -Calculus. Technical Report, University of Berne (2008).
- [AF $\infty$ ] ALBERUCCI L., FACCHINI A.: On Modal  $\mu$ -calculus and Gödel-Löb Logic. To appear in *Studia Logica*.
- [AK $\infty$ ] ALBERUCCI L., KRÄHENBÜHL J.: Justifying Induction on Modal  $\mu$ -Formulae. In Preparation.
- [Arn99] ARNOLD A.: The  $\mu$ -Calculus Alternation-Depth Hierarchy is Strict on Binary Trees. *ITA* 33(4/5): 329-340 (1999)
- [vBe06] VAN BENTHEM, J.: Modal Frame Correspondences and Fixed-Points. *Studia Logica* 83: 133–155 (2006).
- [BRV01] BLACKBURN P., DE RIJKE M., VENEMA Y.: *Modal Logic*. Cambridge University Press (2001)
- [Bra98<sup>1</sup>] BRADFIELD, J.: The Modal  $\mu$ -calculus Alternation Hierarchy is Strict. *Theoretical Computer Science* 195(2): 133–153 (1998).
- [Bra98<sup>2</sup>] BRADFIELD, J.: Simplifying the Modal  $\mu$ -Calculus Alternation Hierarchy. *STACS* 1998: 39-49 (1998)
- [BS01] BRADFIELD, J., STIRLING C.: Modal logic and mu-calculi. In *Handbook of Process Algebra*, edited J. Bergstra, A. Ponse and S. Smolka, 293-332. Elsevier, North-Holland (2001)
- [tCFL07] TEN CATE B., FONTAINE G., LITAK T.: Some Modal Aspects of XPath. *Methods for Modalities* (2007).
- [DL $\infty$ ] D’AGOSTINO G., LENZI G.: On the  $\mu$ -calculus over Transitive and Finite Transitive Frames. Submitted (2008).
- [DO $\infty$ ] DAWAR A., OTTO M.: Modal Characterisation Theorems over Special Classes of Frames. In preparation (2008).
- [EJ91] EMERSON E.A., JUTLA C.S.: Tree Automata,  $\mu$ -Calculus and Determinacy (Extended Abstract). *FOCS* 1991: 368–377 (1991)
- [GTW02] GRÄDEL E., THOMAS W., WILKE T. (Eds.): *Automata, Logics, and Infinite Games*. LNCS 2500, Springer-Verlag Berlin Heidelberg (2002)
- [HC96] HUGHES G.E., CRESSWELL M.J.: *A New Introduction to Modal Logic*. Routledge, London (1996)
- [Koz83] KOZEN D.: Results on the propositional mu-calculus. *Theoretical Computer Science* 27: 333–354 (1983)
- [Koz88] KOZEN D.: A finite model theorem for the propositional mu-calculus. *Studia Logica* 47(3):233-241 (1988)
- [Kir02] KIRSTEN D.: Alternating Tree Automata and Parity Games, in [GTW02]: 157–167.
- [Len96] LENZI G.: A Hierarchy Theorem for the  $\mu$ -Calculus. *ICALP* 1996: 87-97 (1996)
- [Len06] LENZI G.: The Transitive  $\mu$ -Calculus is Büchi-definable. *WSEAS Transactions on Mathematics* 5(9): 1021–1026 (2006)
- [Len08] LENZI G.: Personal Communication, February 26th 2008.
- [SB69] SCOTT D., DE BAKKER, J.: A theory of programs. Unpublished manuscript, IBM, Vienna (1969)
- [Sti01] STIRLING C.: *Modal and Temporal Properties of Processes*. Springer-Verlag Berlin Heidelberg (2001)

- [ES89] STREETT R.S., EMERSON E.A.: An automata theoretic decision procedure for the propositional  $\mu$ -calculus. *Information and Computation* 81(3): 249–264, 1989
- [Maz02] MAZALA R.: Infinite Games. in [GTW02], pp. 23–38
- [Mos91] MOSTOWSKI A.W.: Games with forbidden position. Technical Report 78, University of Gdansk (1991)
- [Tar55] TARSKI A.: A lattice-theoretical fixpoint theorem and its applications. *Pacific Journal of Mathematics* 5(2): 285–309 (1955)
- [Vis96] VISSER, A.: Uniform interpolation and layered bisimulation. In: *Gödel '96 (Brno, 1996)*. Lectures Notes in Logic, 6 Berlin, Springer (1996): 139–164.
- [Vis05] VISSER, A.: Löb's Logic Meets the  $\mu$ -Calculus. In: MIDDELDORP A., VAN OOSTROM V., VAN RAAMSDONK F., DE VRIJER R. (Eds.): *Processes, Terms and Cycles, Steps on the Road to Infinity, Essays Dedicated to Jan Willem Klop on the Occasion of His 60th Birthday*. Berlin, Springer (2005): 14–25.
- [Wal02] WALUKIEWICZ I.: Monadic second-order logic on tree-like structures. *Theoretical Computer Science* 275(1–2): 311–346 (2002).
- [Wil01] WILKE T.: Alternating Tree Automata, Parity Games, and Modal  $\mu$ -Calculus. *Bul. Belg. Math. Soc.* 8(2): 359–391 (2001)